
ADDITIONAL COPIES
OF THIS PUBLICATION MAY BE PROCURED FROM
THE SUPERINTENDENT OF DOCUMENTS
GOVERNMENT PRINTING OFFICE
WASHINGTON, D. C.
AT
10 CENTS PER COPY
SUBSCRIPTION PRICE, \$3 PER YEAR
△

JOURNAL OF AGRICULTURAL RESEARCH

DEPARTMENT OF AGRICULTURE

VOL. V

WASHINGTON, D. C., MARCH 20, 1916

NO. 25.

INHERITANCE OF FERTILITY IN SWINE¹

[PRELIMINARY PAPER]

By EDWARD N. WENTWORTH, *Professor of Animal Breeding*, and C. E. AUBEL, *Fellow in Animal Breeding, Kansas Agricultural Experiment Station*

INTRODUCTION

Mendelian inheritance applies almost without exception to the transmission of qualitative characters. Quantitative traits, on the other hand, are susceptible only to a generalized treatment from this viewpoint, and few investigators have attacked the problem. Size inheritance in animals has been dealt with by Castle and Phillips (2)², Goldschmidt (7), MacDowell (10), Phillips (19, 20), and Punnett and Bailey (21), while Detlefsen (3) has treated the inheritance of certain skeletal characters. Pearl (15) discovered an arbitrary division point of 30 eggs in the winter laying period of hens, for which inheritance apparently depends on two factors, one of which follows an ordinary Mendelian, and the other a sex-linked scheme. These determiners provide the nearest to units of inheritance that have yet been isolated in quantitative studies.

Because of the fact that fecundity deviates only by discrete units, the litter size in swine provides peculiarly favorable material for studying quantitative inheritance. An analysis of this material has already been attempted from the biometric viewpoint. Rommel and Phillips (24) correlated the size of litters in which dams and daughters were farrowed and found a correlation coefficient of 0.0601 ± 0.0086 . They conclude from this result that there is an actual positive correlation between the size of litters of two successive generations, believing that size of litter is a character transmitted from mother to daughter. They recognize the smallness of the coefficient, but believe the indications of inheritance are large enough to provide a basis for selection. In studying fertility inheritance Pearson and Lee (18) obtained practically similar coefficients with the human race and the thoroughbred horse. The range of correlation was 0.0418 to 0.213; hence, they conclude that fertility is certainly and markedly inherited.

¹ Paper No. 1 from the Laboratory of Animal Technology, Kansas Experiment Station.

² Reference is made by number to "Literature cited," p. 1159-1163.

Rommel and Phillips (24) studied the inheritance only through the female line, taking no account of a possible influence of the male. George (6) correlated the size of litter with that of the paternal and maternal grandams, respectively. Only 296 litters were involved in his populations; hence, his probable errors were large. But in the dam and daughter comparisons he approximated very closely the result obtained by Rommel and Phillips. His four coefficients follow:

Daughter and dam.....	0.0615±0.0390
Dam and grandam.....	.1147±.0343
Daughter and maternal grandam.....	.0025±.0392
Daughter and paternal grandam.....	.0508±.0392

None of these correlations are three times as large as their probable error; hence, none are really significant.

Simpson (25) approached the problem from a Mendelian standpoint by crossing a wild German Schwarzwald boar to a young Tamworth sow. The Schwarzwald normally averages 4 pigs to the litter, the Tamworth about 11. The particular sow used was farrowed in a litter of 12 pigs, and to the stint of the wild boar farrowed 9 pigs. In the F_1 generation three females were bred, one to a litter mate and the other two to sires unnamed. The first sow produced 4 pigs, the others 4 and 6, respectively, all in their first litters. The sow producing the 6-pig brood was later served by a pure Schwarzwald boar and farrowed 7 pigs, being apparently constant for that degree of fertility. One of the sows from the brood of 6 gave birth to 12 pigs when mated to a pure Tamworth male. The evidence for a segregation of fecundity factors seems fairly clear, although the numbers are small.

NONGENETIC FACTORS AFFECTING FERTILITY

External factors play a great part in the realization of the inborn hereditary capacity for reproduction. Marshall (12, 13) discusses at length the relation between season and productivity, while the sterility of wild animals in captivity or of domestic animals transferred to vastly different altitudes is proverbial. Marshall and Evvard (4, 5) have both studied the effect of "flushing" in sheep, and Evvard has conducted some very exhaustive investigations into the relation of the various compounds of nutrition to litter size in swine. Using the rate of gain at breeding time in gilts¹ as an indication of the state of nutrition, Evvard has found as much as an average difference of two pigs per litter in favor of the best gainers in each experimental lot, when compared with the poorest gainers. Protein added to a nitrogen-deficient ration (corn alone) produced a marked rise in the fertility of gilts and a medium rise in the fertility of older sows.

Many stockmen believe that overfatness diminishes fecundity. [¶]Here may be both a physical obstruction of the reproductive organs due to

¹ A gilt is a young sow intended for breeding purposes. The term is usually applied only until the first litter is produced, although it is sometimes extended throughout the suckling period.

fat and an adipose degeneration of the sex glands. Whether these are really causes of decreased fertility is doubtful, since the best evidence shows them to be symptoms of reproductive derangement.

Overfatness occurs frequently as a result of disturbances in the metabolism, due to loss of secretion from several of the ductless glands, the sex gland being here included. Castrating or spaying are known to promote obesity; hence, it is quite reasonable to assume that if testicular or ovarian derangement first occurs, then fat deposition will follow. Overfatness would thus merely indicate and not initiate reduced fecundity.

Market hog raisers usually believe that pure-bred hogs are deteriorating in prolificacy, in line with the common idea that inbreeding ultimately results in barrenness. Bitting, in 1898 (1), investigated the average size of the first 200 litters and the last 200 litters recorded at that time in the herdbooks of the Berkshire, Ohio Poland-China, Standard Poland-China, and Improved Chester White registry associations and found that during the period in which registration had taken place the Berkshires had decreased 0.19 pig per litter, the Poland-China had increased 0.225 pig, and the Chester White had increased 0.1 pig. Rommel (22) investigated the same point for a period of 20 years in books of the American and Ohio Poland-China associations, comparing the average size of litter for the first 5 years with the average for the last 5. The increase was 0.62 pig per litter among the American Poland-Chinas and 0.43 pig per litter in the Ohio strain. A similar study by Rommel (22) on the Duroc-Jersey covering over 15 years showed an increase of 0.57 pig. The changes which have occurred here are manifestly opposed to the idea that purity of blood lines diminishes fertility. On the other hand, the purity of blood can not be credited with the increase, since a constant selection for large litters has taken place, although an increased homozygosis for prolificacy might come about gradually with years of such mass selection as ordinary stock breeding involves.

Hammond (8) has shown that ova may be lost either before or after fertilization; and, still more important, he has discovered that a relatively high percentage may atrophy during the earlier stages of embryonic growth. Lewis (9) indicated that there may be morphological interferences with reproduction, so that fertility may be decreased. He found that the sperm cells of the boar are practically all dead after being in the uterus for 48 hours, which would, of course, result in a reduced fertility. Lewis's results on the viability of sperm differ from those of Dührssen (11), who observed living sperms in the Fallopian tubes of a woman patient three weeks after copulation had taken place. The importance of this question is probably confined to individual cases.

Certain relatively extraneous characters are popularly supposed to be correlated with high fertility. Many farmers believe that "big type" or "cold blooded" hogs farrow larger litters than "hot blooded," or

that "Spotted Poland-Chinas" are far more fecund than ordinary strains. Swine judges commonly consider long-bodied sows more prolific than their chubbier mates. A comparison of 1,000 litters of "large type" Poland-Chinas with 1,100 litters of "small type" showed no significant difference in fertility. The mean for the "large type" was 7.854 ± 0.0456 , and for the "small type" was 7.896 ± 0.0436 . Furthermore, the standard deviation of the two groups was almost exactly the same, being 2.142 ± 0.0323 for the former and 2.146 ± 0.0309 for the latter. The writers have never seen more than isolated instances brought forward to confirm the popular ideas on this subject and feel that the bulk of such beliefs have resulted from mere advertising schemes.

Breed certainly has its influence. Bitting (1) has averaged the litter sizes for 400 Berkshires, 1,086 Poland-Chinas, and 600 Chester Whites, with the following results:

Berkshire.....	8.22 pigs per litter.
Poland-China.....	7.45 pigs per litter.
Chester White.....	8.96 pigs per litter.

Surface (26) computed the means and standard deviations in the 54,515 litters of Poland-Chinas and the 21,652 litters of Duroc-Jerseys studied by Rommel (22). His constants follow:

	Mean.	Standard deviation.
Poland-China.....	7.435 ± 0.01	2.038 ± 0.013
Duroc-Jersey.....	$9.337 \pm .021$	$2.427 \pm .016$

The large numbers here involved undoubtedly prove that real breed differences in fertility exist.

Pearl (15, 16) found the number of mammae to be correlated positively with the number at a birth when different species are compared, but the coefficient is very low within the species. Parker and Bullard (14) correlated the same characters in 1,000 litters of swine and obtained a coefficient of 0.0035 ± 0.0124 . The senior author¹ treated the same point in 170 litters of which he had made genetic studies and obtained a coefficient of -0.0059 ± 0.0517 .

These figures certainly demonstrate that the number of mammae in swine is not related to fertility; in fact, nothing so far discussed presents reliable external characters on which fertility selections can be made. Apparently fecundity has as profound a genetic as physiologic basis.

VALUE OF HERDBOOK DATA

There is now on record an immense mass of data relating to fertility inheritance in swine, in the volumes of the different breed registry associations. In addition to the name and number of the animal, its parents, breeder, etc., the size of litter in which it was farrowed is usually stated.

¹ Unpublished data.

This furnishes opportunity to link together any desired number of generations.

In treating such data, the degree of confidence which can be placed in the figure for litter size must be considered. Its accuracy depends on the carefulness and honesty of the breeder, the accuracy of the clerks in the registry office, and the freedom from typographical errors in the printing of the volume. The matter of personal integrity can be accepted to a high degree, for fortunately the majority of breeders are quite reliable. Whenever falsification wittingly occurs, the tendency is to raise the number per litter; but, owing to the publicity involved in pure-bred breeding as well as the personality invested in breeding animals due to the registry systems, it is doubtful if litter sizes are ever exaggerated by more than one or two pigs.

Investigations in color discrepancies, mistakes in parentage, etc., have shown that about 2 per cent of errors are involved in the work of registry-office clerks and in printing. Some associations are more careful than others, but, of course, none are absolutely free from errors. Unfortunately swine books show a relatively greater number of mistakes than do those published by breeders of some of the other classes of live stock.

However, assuming, as has been done, that the bulk of the records can be accepted, there still remains a question as to their genetic value. It is doubtful whether a sow will ever exceed her hereditary possibilities in number per litter, but there are many forces that may cause her to fall short of that number. Lack of proper nutrition, failure to have all ova released or fertilized, loss of ova, atrophy of fertilized ova or embryos, and disease may all operate against the complete realization of the hereditary make-up. The age at which a sow farrows, the number of litters she has per year, and certain other environmental conditions may also reduce the litter size. It is interesting to observe that this source of error operates in a compensating direction to that of record falsification, when such exists, and in the end the two may counterbalance, although these physiological and pathological factors operate more often than does the misrepresentation of litter numbers.

After admitting all of these sources of error, but hoping that enough records are made under natural conditions to give the figures an investigational value, there still remains the big question of the geneticist, Does the somatic expression of the character indicate the germinal (zygotic) condition of the individual? In other words, Does the fact that a pig is farrowed in a litter of eight indicate that it will transmit a tendency to produce litters of eight? The answer very evidently is No, and the greater the degree of outcrossing in the ancestral lines, the less reliable an index of heredity the size of litter is. Yet it is the only single index obtainable in the study of herdbook records; so for the present it will have to be accepted for what it is worth.

ADVANTAGES OF LITTER SIZE INHERITANCE STUDIES

Accepting the figures for litter size as reasonably representative of the hereditary constitution, there are a number of reasons that make them desirable material for inheritance studies. The most important of these is the fact that the male mated to a female probably does not affect the number at a birth. Instead the size of litters a sow produces represents the segregation of the tendencies transmitted to her by her father and mother. Suppose a sow produces a litter of four pigs and is herself from a litter of seven, the seven does not determine in any way the four, but instead the segregation of some tendency transmitted by her sire or dam is represented. The only check available on this tendency in her sire is the size of litter in which he was farrowed, while the same holds for the dam, except that her own breeding performance may give an additional idea.

METHOD OF RECORDING THE DATA

The data on the animals studied were recorded as follows, the figures representing the size of litters in which the individuals were farrowed:

		Grandsire
		4
Animal	Dam	Grandam
4	7	9

The size of litters produced by sows whose sires came from litters of four and whose dams came from litters of seven should give an idea (through the variations recorded) of the hereditary factors involved. It is admissible that all grandams or all grandsires farrowed in the same size of litters may be different in hereditary make-up, but there should be enough individuals alike to make the frequency curves at least suggestive. For convenience, the grandparental generation will be lettered "P," the parental generation " F_1 ," and the filial generation " F_2 ," although it is clearly to be understood that this notation does not have the regular Mendelian significance.

DEVIATIONS PER GENERATION

The mean size of 1,770 litters in the P generation was 7.84 ± 0.3494 . The standard deviation was 2.18 ± 0.2461 . This gives a coefficient of variability of 27.80 for this generation.

The mean size of 385 litters for the F_1 generation was 7.82 ± 0.4897 . The corresponding standard deviation was 2.16 ± 0.3462 . The coefficient of variability here involved was 27.60, practically the same as that of the grandparental generation.

The mean size of 885 litters in the F_2 generation was 7.91 ± 0.4965 , while the deviation was 2.19 ± 0.3511 , giving a coefficient of variability of 27.55. (See Table I.)

TABLE I.—Deviation in size of litters in swine

Generation.	Number of litters.	Mean.	Standard deviation.	Coefficient of variability.
P.....	1,770	7.84 ± 0.3494	2.18 ± 0.2461	27.80
F ₁	885	7.82 ± .4897	2.16 ± .3462	27.60
F ₂	885	7.91 ± .4905	2.19 ± .3511	27.55

The mean litter size is quite constant from generation to generation, and furthermore quite close to that obtained by Surface (26) for the breed in general. If anything of Mendelism is involved here, it is not revealed by this method of treatment, for the standard deviation is so nearly the same for each of the generations involved as to give no hint of segregation. In fact, the coefficients of variability would indicate a slowly increasing degree of homozygosis.

Two interpretations may be placed on these figures. The animals studied are either practically constant from a zygotic standpoint, and the variations in litter size are due to environmental treatment, or else there is so much heterozygosis present in the grandparents that the parents are as much F₂ as F₁ in hereditary make-up. For the present the writers are going to use the latter interpretation, as there is no evidence at hand to support a belief in the former.

TABLE II.—Deviation in litter size of the offspring from the parental generation in swine

BOAR 1						
Size of litter of parents.		Number of matings.	F ₁ generation.		F ₂ generation.	
Boar.	Sow.		Mean.	Standard deviation.	Mean.	Standard deviation.
1	9	1	5	0	9	0
1	4	1	2	0	9	0
BOAR 2						
2	5	1	4	0	6	0
2	6	2	8 ± 1.43	3 ± 1.0117	7.5 ± 0.239	.5 ± 0.1686
2	8	2	6.5 ± .717	1.5 ± .1737	4 ± .479	1 ± .3372
2	9	1	6	0	6	0
BOAR 3						
3	4	3	7 ± 0.171	1.41 ± 0.3433	9 ± 1.704	4.32 ± 1.194
3	6	2	6.5 ± .239	.5 ± .1686	8 ± .95	2 ± .674
3	7	4	7.25 ± 1.6218	4.8 ± 1.1502	7.75 ± .8869	2.63 ± .627
3	11	1	6	0	8	0
3	10	2	10 ± .95	2 ± .674	11	0
3	14	1	11	0	6	0

TABLE II.—Deviation in litter size of the offspring from the parental generation in swine—Continued

BOAR 4

Size of litter of parents.		Number of matings.	F ₁ generation.		F ₂ generation.	
Boar.	Sow.		Mean.	Standard deviation.	Mean.	Standard deviation.
4	3	1	8	0	5	0
4	4	3	7.66 ± 1.032	2.62 ± 0.723	8 ± 0.631	1.63 ± 0.442
4	5	5	7.2 ± .9951	3.29 ± .7022	7.6 ± .5202	1.72 ± .3671
4	6	7	8 ± .2887	1.13 ± .2037	7.14 ± .2836	1.11 ± .2001
4	7	8	8.5 ± .6863	2.42 ± .408	5.87 ± .2726	1.14 ± .1922
4	8	6	7.83 ± .5390	1.95 ± .3801	6.83 ± .7712	2.79 ± .5438
4	9	6	7.16 ± .8127	2.94 ± .5731	7.33 ± .4008	1.45 ± .2826
4	10	4	7.25 ± .5092	1.51 ± .3618	6.75 ± .3844	1.14 ± .2072
4	12	1	7	0	12	0

BOAR 5

5	3	1	6	0	12	0
5	4	4	5.75 ± 0.6407	1.92 ± 0.4543	6.5 ± 0.3743	1.11 ± 0.2654
5	5	3	7 ± .3195	.81 ± .2239	6.66 ± .6666	1.69 ± .4671
5	6	14	8 ± .3228	1.79 ± .2282	7.21 ± .1531	.85 ± .1083
5	7	12	7.91 ± .4404	2.29 ± .358	7.66 ± .3099	1.59 ± .2193
5	8	16	7.37 ± .3979	2.36 ± .2817	7.56 ± .408	2.42 ± .2889
5	9	9	8.11 ± .3844	1.71 ± .272	8.22 ± .5866	2.61 ± .415
5	10	6	9.33 ± .5943	2.15 ± .4191	8.66 ± .3759	1.56 ± .2651
5	11	3	6.66 ± .4891	1.24 ± .3427	8.33 ± .3707	.94 ± .2598
5	12	2	4	0	9	0

BOAR 6

6	2	1	7	0	8	0
6	3	3	7.66 ± 0.8283	2.1 ± 0.5805	3.33 ± 0.6243	1.58 ± 0.4367
6	4	3	4.66 ± 1.1161	2.83 ± .7822	9.33 ± .4391	1.24 ± .3426
6	5	6	8.6 ± .6041	2 ± .4267	6.83 ± .4533	1.64 ± .3197
6	6	11	6.72 ± .4604	2.26 ± .3251	8.9 ± .4625	2.27 ± .3203
6	7	18	7.05 ± .2672	1.68 ± .1888	8.5 ± .252	1.86 ± .178
6	8	24	7.58 ± .3255	2.36 ± .2300	7.46 ± .1862	1.35 ± .1313
6	9	15	8.4 ± .0040	2.74 ± .0311	8 ± .047	2.7 ± .0314
6	10	8	6.62 ± .4657	1.71 ± .2885	7.87 ± .6075	2.54 ± .4283
6	11	5	9 ± .9151	2.32 ± .6414	5 ± .6922	2.39 ± .4887
6	12	3	7.66 ± .1853	.47 ± .1297	8.33 ± 1.136	2.88 ± .790

BOAR 7

7	3	1	6	0	10	0
7	4	4	7.5 ± 0.8431	2.5 ± 0.5802	8.5 ± 0.1631	5 ± 0.119
7	5	7	7.71 ± .4202	1.62 ± .292	7.85 ± .6100	2.39 ± .431
7	6	17	7.25 ± .2177	1.33 ± .1541	7.58 ± .239	1.46 ± .1691
7	7	19	7.52 ± .3023	1.95 ± .2135	7.42 ± .3178	2.05 ± .2244
7	8	26	8.03 ± .3637	1.99 ± .1843	7.8 ± .2545	1.92 ± .1778
7	9	41	9.29 ± .1958	1.83 ± .1363	8.82 ± .2077	1.94 ± .1445
7	10	8	8.37 ± .5333	2.23 ± .3760	6.62 ± .4616	1.93 ± .3254
7	11	16	8.5 ± .2506	1.54 ± .1838	8.43 ± .4502	2.76 ± .3187
7	12	6	8.66 ± .5943	2.15 ± .4191	6.5 ± .6109	2.21 ± .4303
7	13	1	10	0	4	0

TABLE II.—Deviation in litter size of the offspring from the parental generation in swine—Continued

			BOAR 8			
Size of litter of parents.			F ₁ generation.		F ₂ generation.	
Boar.	Sow.	Number of matings.	Mean.	Standard deviation.	Mean.	Standard deviation.
8	3	7	7.28±0.1124	0.44±0.0793	7 ±0.3193	1.25±0.2254
8	4	8	8.75±.6721	2.81±.4738	7.75±.5572	2.33±.3928
8	5	11	7.33±.2634	1.91±.1861	8.27±.6377	3.13±.4591
8	6	24	6.91±.2965	2.15±.2094	7.66±.3062	2.22±.2161
8	7	29	7.86±.2394	1.91±.1692	7.82±.277	2.21±.1958
8	8	43	6.58±.2519	2.45±.1783	7.23±.2076	2.02±.1469
8	9	34	8.14±.292	2.52±.2124	8.17±.2294	1.98±.1669
8	10	10	6.7 ±.2219	1.04±.1565	8.4 ±.1462	5.37±.8683
8	11	10	7.8 ±1.024	4.8 ±.7225	8.6 ±.512	2.4 ±.3612
8	12	3	9 ±1.4061	3.55±.9895	9.66±.8415	2.04±.5639
8	13	1	9	0	10	0
BOAR 9						
9	2	1	5	0	9	0
9	3	1	5	0	7	0
9	4	6	7.83±0.4616	1.67±0.3254	7.66±0.4616	2.05±0.3996
9	5	12	7.5 ±.3898	2 ±.2758	6.91±.4482	2.3 ±.3172
9	6	7	6.71±.4444	1.74±.3138	7.14±.6065	2.35±.4238
9	7	32	7.59±.2232	1.87±.1576	8.53±.2495	2.09±.1762
9	8	26	7 ±.2663	2.01±.1861	7.5 ±.3074	2.32±.2148
9	9	35	7 ±.3172	2.95±.2377	8.71±.2158	2.10±.1693
9	10	14	8.66±.4060	2.33±.2874	7.53±.3468	1.99±.2453
9	11	8	7.75±.409	1.71±.2883	8.5 ±.526	2.2 ±.3709
9	12	4	8 ±.1686	.5 ±.119	9.25±.3709	1.1 ±.263
9	13	1	7	0	8	0
9	15	1	6	0	8	0
BOAR 10						
10	1	1	8	0	14	0
10	3	2	9 ±0.4784	1 ±0.3372	7.5 ±1.674	3.5 ±1.1803
10	4	2	9	0	7.5 ±1.674	3.5 ±1.1803
10	5	4	7 ±.4755	1.41±.3372	7.25±.4957	1.47±.3516
10	6	6	7.66±.34	1.23±.2397	8 ±.3897	1.41±.2749
10	7	16	7.03±.3591	2.13±.2542	8.18±.4991	2.96±.3533
10	8	24	8.29±.2978	2.16±.2105	8.26±.3787	2.69±.2676
10	9	11	8.9 ±.6153	3.02±.4343	8.63±.3586	1.76±.2531
10	10	14	6.78±.4128	2.20±.2019	7.57±.2777	1.54±.1963
10	11	4	8.25±.435	1.29±.3085	9.5 ±.5508	1.66±.397
10	12	6	8.5 ±.525	1.9 ±.370	8.63±.5888	2.13±.4151
10	14	1	10	0	8	0
BOAR 11						
11	3	1	9	0	10	0
11	4	1	6	0	9	0
11	5	5	6.8 ±0.0574	1.9 ±0.0405	8.2 ±0.5111	1.69 ±0.3607
11	6	6	8 ±.5749	2.08±.4054	6.33±.4644	1.68 ±.3275
11	7	12	7.33±.3004	1.54±.2124	7.85±.3804	2.007±.2705
11	8	4	8.5 ±.1686	.5 ±.1190	8.75±1.2107	3.59 ±.8586
11	9	6	8.83±.6053	2.19±.4269	8.16±.4809	1.74 ±.3391
11	10	6	7.5 ±.3434	1.25±.2422	8.66±.4046	1.69 ±.3277
11	13	1	6	0	12	0
11	15	1	9	0	9	0

TABLE II.—Deviation in litter size of the offspring from the parental generation in swine—Continued

BOAR 12						
Size of litter of parents.		Number of matings.	F ₁ generation.		F ₂ generation.	
Boar.	Sow.		Mean.	Standard deviation.	Mean.	Standard deviation.
12	2	1	9	0	7	0
12	4	2	9.5 ± 0.238	.5 ± 0.118	8.5 ± 0.238	.5 ± 0.118
12	5	2	4 ± .956	2 ± .6745	7	0
12	6	1	8	0	10	0
12	7	5	7.6 ± .554	1.85 ± .3001	9.2 ± .399	1.32 ± .282
12	8	6	8.82 ± .384	1.39 ± .27	8.5 ± .671	2.4 ± .473
12	9	6	6.5 ± .591	2.14 ± .414	7.83 ± .387	1.4 ± .272
12	10	1	11	0	10	0
12	11	2	11	0	8.5 ± .717	1.5 ± .505
12	12	1	10	0	12	0
12	13	2	7	0	9.5 ± .717	1.5 ± .505
BOAR 13						
13	6	3	8.66 ± 0.493	1.25 ± 0.345	8.33 ± 0.185	0.47 ± 0.129
13	8	2	9.5 ± .717	1.5 ± .505	7 ± .956	2 ± .6745
13	9	2	11.5 ± .717	1.5 ± .505	9 ± .956	2 ± .6745
13	11	1	10	0	6	0
13	12	2	7.5 ± .168	.5 ± .243	10 ± .479	1 ± .337
13	13	5	9.6 ± .526	1.74 ± .371	10 ± .956	2 ± .6745
BOAR 14						
14	8	2	9	0	10.5 ± 0.239	0.25 ± 0.1686
14	9	1	9	0	11	0
14	12	1	7	0	3	0
BOAR 15						
15	8	1	12	0	8	0

INDIVIDUAL EVIDENCES OF SEGREGATION

Table II is produced by treating the litter size as a detailed character and comparing the parental generation with offspring. The average of the F₁ deviations is 1.87 ± 0.0549 , while the F₂ mean deviation is 1.92 ± 0.0582 . The probable errors make these two constants overlap, so that the individual treatment when lumped seems no more significant than when the deviations per generation are considered. Yet many individual evidences of segregation exist, and many times the F₂ generation from a particular cross is so small in numbers that only a fragmentary view of the segregable possibilities is obtained.

While it is possible that 90 per cent of the litter sizes in these tables do not represent the exact genetic constitution, yet it is probable that in general the greater the disparity in litter sizes between the two animals in the P generation, the greater will be the expected deviations in the F₂, and the smaller the deviations in the F₁ generation. The following results, Table III, are produced by tabulating the averages of the deviations on this basis.

TABLE III.—Average deviations in litter size in the F₁ and F₂ generations of swine

Difference in number of pigs in the two P litters.	0	1	2	3	4	5	6	7	8
F ₁ deviations	2.13	1.89	1.93	2.04	2.19	1.82	1.32	1.12	0.5
F ₂ deviations	1.91	1.84	2.16	2.10	2.16	1.71	1.72	1.98	.5

A calculation of the probable errors involved in this table shows that only the difference between the F₁ and F₂ deviations where the disparity in litter size is seven pigs is large enough to be mathematically significant. The difference when the parents vary from each other by two pigs and by six pigs is on the border line between significance and nonsignificance, but the five other columns are distinctly unenlightening. Yet, if the difference of two pigs is barred, the results are what might be expected.

One criticism against the preceding method of treatment is thoroughly valid. If swine fertility depends on only one or two genetic factors, it is obvious that the point at which the difference between the two parents occurs is more important than the degree of difference. For example, if there is a physiological division point between two hereditary factors at six pigs, then a difference of two or even of four below six pigs might not be significant, while a difference of one more or one less in a litter of six or seven pigs would be thoroughly significant. An examination of the data from this point of view is now in progress, but it is probable that the key to the situation will only be discovered by breeding experiments.

CURVES OF LITTER FREQUENCIES

The distribution of the different sizes of litters in the three generations is given in Table IV.

TABLE IV.—Litter frequencies in swine

Generation.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
P Expected	0.11	1.5	9.8	39	108	216	324	370	324	216	108	30	9.8	1.5	0.11
P Actual	3	9	30	82	124	198	300	302	318	162	91	59	20	6	3
F ₁ Expected	0	0.05	0.75	4.9	19	54	100	162	185	162	108	54	19	4.9	0.75
F ₁ Actual	0	5	14	32	69	122	149	161	149	85	62	25	8	4	2
F ₂ Expected	0	0.05	0.75	4.9	19	54	100	162	185	162	108	54	19	4.9	0.75
F ₂ Actual	0	4	17	32	63	107	154	172	135	95	59	30	11	3	3

Figures 1, 2, 3, and 4 show the curves for the litter frequencies in the three generations and indicate how close the actual numbers of litters come to the binomial curve $(x + y)^n$. It is perhaps incorrect to call the theoretical frequencies recorded in Table IV "expectations," unless it is clearly understood that they are the expectations founded on the nearest binomial. There is nothing in the inheritance to make them true expectations from an experimental standpoint.




FIG. 1.—Curve of litter frequencies in the P generation of swine.

mean for each perfectly valid. The modes of these three curves are as follows:

Curve 1.....	4 pigs per litter.
Curve 2.....	8 pigs per litter.
Curve 3.....	12 pigs per litter.

It is premature to announce that these modes represent centers of deviation for genetic factors, although a casual observation of the individual data makes it seem that this condition may exist. Furthermore, the mode of curve 1 corresponds to the degree of fertility which Simpson states is characteristic of the wild hog, while the mode of curve 3 is very close to that of the Tamworth, the most fecund of domestic breeds. This indicates that the two may represent basic and improved factors for fertility, respectively, while curve 2 represents heterozygous condition.

Before these curves can be accepted as more than merely suggestive a further analysis must be made. There is a significant deviation from expectancy in the right-hand branch of the curve of the total population, which persists even after the separation into three curves. In figure 4 this deficiency is located in the left-hand branch of curve 3, but the minus deviations may just as logically belong in the right-hand branch of curve 2, suggesting that it also may be compounded of two curves dependent on a genetic factor not disclosed thus far.

Paralleling this study some actual matings of swine have been planned and are in progress.

SUMMARY

(1) Fertility in swine offers favorable material for the study of quantitative inheritance, because the units of deviation are discrete.

(2) Biometric studies of litter size with mother and daughter have indicated a small degree of inheritance.

(3) Crosses of breeds having different mean




FIG. 1.—Curve of litter frequencies in the F_1 generation of swine.




FIG. 3.—Curve of litter frequencies in the F_2 generation of swine.

litter sizes have suggested that segregations of fecundity factors may take place.

(4) Numerous nongenetic factors limit the full expression of the inborn possibilities of fertility.

(5) Certain few somatic characters may be correlated either in a physiological or genetic manner with the different degrees of fecundity, but the bulk of characters usually assumed to be so related are probably

entirely independent of it.

(6) Herdbook data on the fertility of swine present sources of error, but the percentage of error is low enough to permit the statistics to be suggestive.

(7) Numerous influences exist which lower the size of litter, which sources of error may operate in a manner compensatory to those just mentioned.

(8) It is questionable whether the size of litter represents the hereditary factors transmitted, but the somatic character was perforce accepted at face value in these studies.

(9) There is no reduction in variability




FIG. 4.—Diagram of the combined litter frequencies for the three generations of swine analyzed into its component curves.

in the litter sizes of the dams as compared with the grandparents or progeny, as would result if there were homozygous differences for fertility in the grandparents. Hence, the fertility deviations are either non-germinal or else the degree of heterozygosis is so great in the grandparents that no increased variability in the F_2 generation is possible. The latter explanation is probably the correct one.

(10) The frequency curves for the 3,540 litters studied make it appear that there are at least three centers of deviation in swine fertility. These centers possibly correspond to genetic factors involved in the inheritance of fecundity.

LITERATURE CITED

- (1) BITTING, A. W.
1898. The fecundity of swine. *In* 10th Ann. Rpt. Ind. Agr. Exp. Sta. 1897, p. 42-46.
- (2) CASTLE, W. E., and PHILLIPS, J. C.
1914. Piebald Rats and Selection; an Experimental Test of the Effectiveness of Selection and of the Theory of Gametic Purity in Mendelian Crosses. 56 p., 3 pl. Washington, D. C. (Carnegie Inst. Washington Pub. 195.) Bibliography, p. 31.
- (3) DEYLEFSEN, J. A.
1914. Genetic Studies on a Cavy Species Cross. 134 p., 10 pl. (1 col.). Washington, D. C. (Carnegie Inst. Washington Pub. 205.) Bibliography, p. 129-132.
- (4) EVVARD, J. M.
1912. Nutrition as a factor in fetal development. *In* Ann. Rpt. Amer. Breeders' Assoc., v. 8, p. 549-560.
- (5) ———
1913. Some factors affecting fetal development. *In* Proc. Iowa Acad. Sci., v. 20, p. 325-330.
- (6) GEORGE, C. R.
1912. The inheritance of fecundity in Poland-China swine. Thesis, University of Ohio.
- (7) GOLDSCHMIDT, Richard.
1913. Zuchtversuche mit Enten I. *In* Ztschr. Indukt. Abstamm. u. Vererbungslehre, Bd. 9, Heft 3, p. 161-191.
- (8) HAMMOND, John.
1914. On some factors controlling fertility in domestic animals. *In* Jour. Agr. Sci., v. 6, pt. 3, p. 263-277, pl. 3.
- (9) LEWIS, L. L.
1911. The vitality of reproductive cells. *Okla. Agr. Exp. Sta. Bul.* 96, 47 p., 7 fig.
- (10) MACDOWELL, E. C., and CASTLE, W. E.
1914. Size Inheritance in Rabbits . . . with a Prefatory Note and Appendix, by W. E. Castle. 55 p., illus. Washington, D. C. (Carnegie Inst. Washington Pub. 196.) Bibliography, p. 47-49.
- (11) McMURRICH, J. P.
1913. Development of the Human Body . . . ed. 4, 495 p., 285 fig. Philadelphia.
Cites (p. 35) case reported by Dührssen.
- (12) MARSHALL, F. H. A.
1908. Fertility in Scottish sheep. *In* Trans. Highland and Agr. Soc. Scot., s. 5, v. 20, p. 139-151.
- (13) ———
1910. The Physiology of Reproduction . . . with a Preface by E. A. Schäfer . . . and Contributions by William Cramer . . . and James Lochhead. 706 p. illus. New York, London.
- (14) PARKER, G. H., and BULLARD, C.
1913. On the size of litters and the number of nipples in swine. *Proc. Amer. Acad. Arts and Sci.*, v. 40, no. 7, p. 397-426. Bibliography, p. 413.
- (15) PEARL, Raymond.
1912. The mode of inheritance of fecundity in the domestic fowl. *In* Jour. Exp. Zool., v. 13, no. 2, p. 153-268, 2 fig. Literature cited, p. 266-268.

- (16) PEARL, Raymond.
1913. On the correlation between the number of mammae of the dam and size of litter in mammals. I. Interracial correlation. *In Proc. Soc. Exp. Biol. and Med.*, v. 11, no. 1, p. 27-30.
- (17) ———
1913. On the correlation between the number of mammae of the dam and size of litter in mammals. II. Intraracial correlation in swine. *In Proc. Soc. Exp. Biol. and Med.*, v. 11, no. 1, p. 31-32.
- (18) PEARSON, Karl, LEE, Alice, and BRAMLEY-MOORE, Leslie.
1899. Mathematical contributions to the theory of evolution. VI. Genetic (reproductive) selection: Inheritance of fertility in man, and of fecundity in thoroughbred racehorses. *In Phil. Trans. Roy. Soc. London, s. A*, v. 192, p. 257-330.
- (19) PHILLIPS, J. C.
1912. Size inheritance in ducks. *In Jour. Exp. Zool.*, v. 12, no. 3, p. 369-380. Bibliography, p. 380.
- (20) ———
1914. A further study of size inheritance in ducks with observations on the sex ratio of hybrid birds. *In Jour. Exp. Zool.*, v. 16, no. 1, p. 131-148. Bibliography, p. 145.
- (21) PUNNETT, R. C., and BAILEY, P. G.
1914. On inheritance of weight in poultry. *In Jour. Genetics*, v. 4, no. 1, p. 23-39, 2 fig., pl. 4.
- (22) ROMMELL, G. M.
1906. The fecundity of Poland-China and Duroc-Jersey sows. U. S. Dept. Agr. Bur. Anim. Indus. Circ. 95, 12 p.
- (23) ———
1907. The inheritance of size of litter in Poland China sows. *In Amer. Breeders' Assoc. Rpt.* v. 3, p. 201-208.
- (24) ——— and PHILLIPS, E. F.
1906. Inheritance in the female line of size of litter in Poland China sows. *In Proc. Amer. Phil. Soc.*, v. 45, no. 184, 245-254. Bibliography, p. 254.
- (25) SIMPSON, Q. I.
1912. Fecundity in swine. *In Ann. Rpt. Amer. Breeders' Assoc.*, v. 7, p. 261-266.
- (26) SURFACE, F. M.
1909. Fecundity of swine. *In Biometrika*, v. 6, pt. 4, p. 433-436.

RELATION OF GREEN MANURES TO THE FAILURE OF CERTAIN SEEDLINGS

By E. B. FRED,

*Agricultural Bacteriologist, Agricultural Experiment Station of the University of Wisconsin*¹

INTRODUCTION

In a previous report it has been shown that if green manures are turned under and the soil planted immediately, a decrease in germination may result. For example, a 20-acre field, half in crimson clover (*Trifolium incarnatum*) and half in fallow, was plowed and planted to cotton (*Gossypium* spp.) (17, p. 26).² On the crimson-clover plot the cotton failed almost completely to germinate. Here and there a few crippled seedlings appeared, while on the fallowed plot normal germination occurred. Seed from the same lot was used on both plots. The green manure in some way seriously affected the germination of the cottonseed. Three weeks later the green-manure plot was again seeded to cotton. Germination at this time was perfectly normal. Apparently the harmful factor disappeared during the interval of three weeks.

A more extensive study of the substances affecting seed germination and of the factors involved was deemed advisable. The controlling idea in this investigation was a study of the effect of green manures on the germination of different seeds. In determining the percentage of germination, only those seedlings that appeared above the surface are recorded.

The amount of green manure used was determined from the following calculation: A good crop of clover should yield from 4 to 5 tons of undried green hay per acre. If 1 acre of soil 3 inches deep weighs 1,000,000 pounds, then 1 per cent of green clover is comparable to the amount employed under field conditions. Except in rare cases this amount of green manure was used in all of the laboratory studies. The green plant tissue was cut just before blooms began to form, finely chopped, and mixed thoroughly with Miami silt loam soil from the Experiment Station farm. The soil moisture was maintained at 50 per cent saturation. All tests of germination are recorded in percentages. Photographs were made of the young seedlings two weeks after planting.

EFFECT OF GREEN MANURES ON THE GERMINATION OF VARIOUS SEEDS

Since it has been shown that seeds of different plants vary widely in chemical composition, it is very probable that they will react differently toward green manures. This experiment was planned to test the effect

¹ Published with the permission of the Director of the Wisconsin Agricultural Experiment Station.

² Reference is made by number to "Literature cited," p. 1175-1176.

of decomposing plant tissue on the germination of buckwheat, castor beans, corn, crimson clover, flax, hemp, lupines, mustard, oats, peanuts, soybeans, sunflower, and wheat. The percentage composition of these seeds is given in Table I.

TABLE I.—The percentage composition of various seeds (11, 20)

Name.	Fat.	Crude protein.	Nitrogen-free extract.	Crude fiber.	Ash.
Castor bean (<i>Ricinus communis</i>)	51.37	18.75	1.5	18.1	3.1
Peanut (<i>Arachis hypogaea</i>)	45	25	18	2 to 5	4.3
Flax (<i>Linum usitatissimum</i>)	33.7	22.6	23.2	7.1	4.3
Hemp (<i>Cannabis sativa</i>)	32.58	18.23	21.06	14.97	4.24
White mustard (<i>Brassica alba</i>)	29.66	27.59	20.83	10.27	4.47
Sunflower (<i>Helianthus annuus</i>)	28.79	16.3	17.28	27.9	3.3
Cotton (<i>Gossypium herbaceum</i>)	20.86	19.69	23.43	21.1	3.8
Soybean (<i>Glycine soja</i>)	17.00	35.00	26.00	5 to 6	4.5
White lupine (<i>Lupinus albus</i>)	6.79	28.78	33.65	11.02	2.99
Oat (<i>Avena sativa</i>)	5.27	10.25	59.68	9.97	3.02
Corn (<i>Zea mays</i>)	4.5	9.5	68.5	2.18	1.6
Buckwheat (<i>Fagopyrum tataricum</i>)	2.68	11.41	58.79	11.44	2.38
Wheat (<i>Triticum sativum</i>)	1.65	10.93	70.01	2.12	1.92

The seeds are grouped according to fat content; those richest in fat are given first. The marked difference in the chemical composition of various seeds is very noticeable. For instance, castor beans contain more than 50 per cent of fat, while oats contain less than 2 per cent.

According to Nobbe (16, p. 173), seeds rich in oil require more oxygen for germination than starch seeds. In Tables II, III, and IV data are presented concerning the effect of green manures on various seeds. In every case the seeds were tested under identical conditions. The figures of Table II show the effect of 1 per cent of green clover on the germination of buckwheat, corn, hemp, lupine, and sunflower.

TABLE II.—Effect of green clover on the germination of various seeds

No.	Seed.	Treatment.	Germination.		
			1 week.	2 weeks.	Relative.
			Per cent.	Per cent.	Per cent.
1	Buckwheat	None	75	90	100
2	do	1 per cent clover	90	90	100
3	Corn	None	100	100	100
4	do	1 per cent clover	95	100	100
5	Hemp	None	95	95	68
6	do	1 per cent clover	65	65	100
7	Lupine	None	75	80	100
8	do	1 per cent clover	60	60	75
9	Mustard	None	95	95	58
10	do	1 per cent clover	55	55	100
11	Sunflower	None	90	90	100
12	do	1 per cent clover	90	90	100

The average percentage of germination in duplicate pots, after one and two weeks, is recorded in Table II. The last column gives the relation between the treated and untreated seeds. A glance at the figures shows clearly that buckwheat, corn, and sunflower were not injured by green manures. On the other hand, hemp and mustard were seriously injured; the latter showed the greatest loss. Lupines are not so sensitive as mustard or hemp toward green manure, although a slight decrease in germination is noted.

As regards fat content, it will be seen that with the exception of sunflower those seeds rich in oil are the most sensitive to green manuring. The very quick germination of sunflower seed may explain their resistance to the injurious factor.

Table III presents data to show the striking difference in behavior of fat and starch seeds toward green manures. A comparison of the injury resulting from the use of green clover and green oats is made.

TABLE III.—*Effect of green clover and oats on the germination of cottonseed and wheat*

No.	Seed.	Treatment.	Germination.			
			1 week.	2 weeks.	3 weeks.	Relative.
			Per cent.	Per cent.	Per cent.	Per cent.
1	Cotton.....	None.....	85	92.5	92.5	100
2do.....	1 per cent of oats....	45	65	65	70
3do.....	1 per cent of clover..	17.5	17.5	17.5	19
4	Wheat.....	None.....	95	100	100	100
5do.....	1 per cent of oats....	85	90	90	90
6do.....	1 per cent of clover..	85	85	85	85

The germination of cotton was seriously injured by the presence of green manures; the green clover was much more harmful than oat tissue. Wheat was little affected by the use of green manure. The data confirm the results of the preceding test—that is, that seeds rich in oil are especially sensitive to green manures. It appears that the percentage of injury depends to a certain degree on the source of the plant tissue. Plate LXXXIII, figure 1, is reproduced from a photograph of cotton seedlings two weeks after planting. In order to make the seedlings more visible, a thin layer of white quartz sand was poured upon the surface of the soil.

With soybeans in place of wheat, this experiment was repeated, as shown in Table IV.

TABLE IV.—Effect of green clover and oats on the germination of cottonseed and soybeans

No.	Seed.	Treatment.	Germination.			
			1 week.	2 weeks.	3 weeks.	Relative.
			<i>Per cent.</i>	<i>Per cent.</i>	<i>Per cent.</i>	<i>Per cent.</i>
1	Cotton	None	95	100	100	100
2	do	1 per cent of oats	35	35	35	35
3	do	1 per cent of clover		10	10	10
4	Soybean	None	100	100	100	100
5	do	1 per cent of oats	40	40	40	40
6	do	1 per cent of clover	30	60	60	60

Here it was again found that the oil seeds are very sensitive to green-manuring. Soybeans are more resistant to this injury than cotton.

As regards the source of the green manure, the results of numerous tests indicate that clover causes a greater loss than oat tissue. An exception to this is found with soybeans (Table IV). No satisfactory explanation has been found for the different action of these two substances. The average of three total-nitrogen analyses shows that clover contains 80.27 per cent of moisture and 4.8 per cent of protein (dry basis). The oats contained 82 per cent of moisture and 3.96 per cent of protein. Chemical analyses fail to disclose any very striking differences between the clover and oat tissue. Indeed, the protein content is nearly the same in both substances. It is possible that the nitrogen of legumes is more available than that of nonlegumes (14). It was noticed repeatedly that clover tissue decomposes more rapidly than oat tissue.

EFFECT OF TIME OF PLANTING AND QUANTITY OF GREEN MANURE ON THE GERMINATION OF COTTON SEED

Ten half-gallon jars were filled with soil and treated as shown in Table V.

TABLE V.—Effect of time of planting and quantity of clover on the germination of cottonseed

No.	Treatment.	Germination.							
		Planted immediately.				Planted two weeks later.			
		1 week.	2 weeks.	3 weeks.	Relative.	1 week.	2 weeks.	3 weeks.	Relative.
		<i>Per ct.</i>	<i>Per ct.</i>	<i>Per ct.</i>	<i>Per ct.</i>	<i>Per ct.</i>	<i>Per ct.</i>	<i>Per ct.</i>	<i>Per ct.</i>
1	None	90	90	90	100	90	95	95	100
2	0.25 per cent of clover	60	60	60	66	90	95	95	100
3	0.5 per cent of clover	50	50	50	55	80	95	95	100
4	1.0 per cent of clover	35	35	35	38	100	100	100	100
5	2.0 per cent of clover					75	85	85	89
6	3.0 per cent of clover					70	85	85	89

From the data of this experiment it is very evident that the serious injury caused by green manures is only temporary. Two weeks after the green manure was turned under, the conditions that affect seed germination disappeared. Aside from the temporary nature of the injurious agent, it will be seen that the percentage of injury is fairly proportionate to the amount of green clover used. In the presence of 0.25 per cent, the rate of germination was decreased 34 per cent, while more than 1 per cent of green manure entirely prevented germination. A comparison of the effect of green manures in different stages of decomposition on cotton germination is shown in Plate LXXXIII, figures 2 and 3.

FIELD EXPERIMENTS WITH GREEN MANURES

Early in the spring of 1914 a series of plot experiments with various seeds was made. For this purpose a good clover sod from the Experiment Station farm, near Madison, Wis., was chosen. This sod was divided into three equal sections: *A*, Clover; *B*, oats; and *C*, unplanted. The sections were subdivided into six plots, as shown in Table VI. Section *A* was allowed to remain in clover, while *B* and *C* were plowed, section *B* planted to oats, and *C* left without any crop. When the oats in section *B* and the clover in section *A* were partly in bloom, the soil was plowed and prepared for planting. One half of each section was planted immediately, the other half 25 days later. It was arranged to study the effect of clover and oat tissue on the germination of cotton, corn, hemp, oats, and soybeans. The same weight of seed was planted in each plot. The results of this series of tests are given in Tables VI and VII.

TABLE VI.—*Effect of green clover on the germination of various seeds*

No.	Seed.	Planted immediately after turning under.				Germination of seed planted 25 days after turning under.	
		With clover.		Unplanted.		With clover.	Unplanted.
		Seed germination.	Weight.	Seed germination.	Weight.		
			<i>Pounds.</i>		<i>Pounds.</i>		
1	Cotton.....	60	91	190	210
2	do.....	71	129	202	218
3	Corn.....	76	21	79	27	68	75
4	Hemp.....	Few.	8	Many.	27	1,050	1,130
5	Oats.....	505	474	Fine.	Fine.
6	Soybean.....	58	4	83	5.5	83	88

TABLE VII.—*Effect of oats on the germination of various seeds*

No.	Seed.	Germination of seed.			
		Planted immediately after turning under.		Planted 25 days after turning under.	
		With oats.	Unplanted.	With oats.	Unplanted.
1	Cotton.....	100	210	134	140
2	do.....	117	218	125	131
3	Corn.....	62	75	72	73
4	Hemp.....	450	1, 130	210	320
5	Oats.....	Many.	Many.	Many.	Many.
6	Soybean.....	35	88	39	40

From these tables it will be seen that green manures seriously injure the germination of cotton, soybeans, and hemp, while corn and oats are not affected. The diminished germination is not confined to clover tissue, but is noted with oats. This effect of the plant tissue on germinating seeds is also observed in the weight of harvest. Unfortunately, because of climatic conditions, the cotton could not be grown to maturity. On adjoining plots, where the green manure was allowed to decompose for 25 days before planting, no injury was observed.

The field data show (1) that green manures largely prevent the germination of certain oil seeds, and (2) that the unfavorable condition is only temporary.

NATURE OF THE INJURIOUS AGENT

There are a number of possible causes that might account for the destructive influence of green manures on seed germination:

First, the green manure greatly increases the number and variety of micro-organisms. The organisms on the plant tissue may be harmful, or conditions proper for the development of harmful organisms may arise.

Second, the large gain in number of organisms, after the addition of green manure, results in a possible accumulation of substances toxic to germination—for example, poisonous by-products of decomposition, as alkali or acid.

Third, the rapid multiplication of micro-organisms, which results in an increased metabolism, causes soil oxygen to be consumed and carbon dioxide to be given off. Such loss in oxygen and gain in carbon dioxide might conceivably retard or prevent germination. If it is assumed that oil seeds require more oxygen for germination than starch seeds, the third supposition should apply particularly to seeds rich in fat (16, p. 173).

EFFECT OF SOIL TYPE

In order to ascertain the relation to soil type of the agent causing a decrease in germination, a series of tests was made. Four soil types were used: Colby silt loam, Miami silt loam, Sparta acid sand, and neutral sand. The results of the first test are given in Table VIII.

TABLE VIII.—*Effect of green manure on the germination of cottonseed*

No.	Soil.	Treatment.	Germination.			Relative.
			1 week.	2 weeks.	3 weeks.	
1	Colby silt loam (acid).....	None.....	Per ct. 90	Per ct. 90	Per ct. 90	Per ct. 100
2	do.....	1 per cent of clover	35	45	50	55
3	Miami silt loam.....	None.....	75	75	75	100
4	do.....	1 per cent of clover.	35	35	35	50
5	Miami silt loam, half sand....	None.....	95	95	95	100
6	do.....	1 per cent of clover.	45	45	45	50
7	Sand.....	None.....	80	80	80	100
8	do.....	1 per cent of clover.	90	90	90	112
9	Sparta acid sand.....	None.....	80	80	85	100
10	do.....	1 per cent of clover	70	70	70	82

For the purpose of securing variation in texture, dilutions with Miami soil and quartz sand were made. From the data obtained, it seems that the property of reducing seed germination is common to both silt loams, but is absent or almost inactive in the sands. Since the relative decrease in germination is approximately the same with Miami or Colby silt loam, it appears that soil reaction is not one of the controlling factors. In neutral or acid sand no decided injury was noted. The results of a second series of tests confirm the above statement. Just why sandy soil should prove less efficient than the loams is not evident from the data, unless it is due to the absence of the injurious factor.

EFFECT OF POSITION OF GREEN MANURE

It was arranged to study the effect on seed germination of plant tissue at different depths. Green clover was added at the rate of 1 per cent. The results secured were as follows: When the green manure was placed in the bottom of the jar, 80 per cent of cotton germinated; in the middle, none germinated; on top, 10 per cent germinated. It is evident that green clover must be in close contact with the seed in order to be effective. This may be shown by wrapping cotton seeds with clover leaves. One or two clover leaves greatly injured seed germination. Plate LXXXIII, figure 4, shows the effect of position of green manure on seed germination.

EFFECT OF INCREASED AERATION

In view of the different action of green manures in compact and open soils, it was decided to make a series of tests under conditions that tend to remove gaseous substances. For this purpose, specially designed jars with openings in their bottoms were employed. By means of a glass tube connected with the bottoms of the jars, air was forced through the soil. In these tests air was allowed to pass through the soil for 20 to 30 minutes every day. A comparison of germination in the aerated and unaerated soils failed to show any difference. Change in soil air did not lessen the injury.

EFFECT OF TEMPERATURE

It is a well-known fact that slight changes in temperature often greatly increase or decrease the growth of micro-organisms. Accordingly a test was made with three variations in temperature.

TABLE IX.—*Effect of temperature on germination of cottonseed*

No.	Treatment.	Temperature.	Germination.		Relative.
			4 days.	8 days.	
		* C.	Per cent.	Per cent.	Per cent.
1	None	25	85	85	100
2	1 per cent of clover	25	55	55	64
3	None	30	95	95	100
4	1 per cent of clover	30	35	35	36
5	None	37	100	100	100
6	1 per cent of clover	37	80	80	80

About 30° C. seems to give the greatest injury; lower or higher temperatures fail to cause so great a decrease in germination.

EFFECT OF CERTAIN DECOMPOSITION PRODUCTS

In the decomposition of plant tissue many substances are liberated—e. g., ammonia and carbon dioxide. The relation of ammonium hydroxid to seed germination has been studied by Bokorny (3; 4, p. 37). He found that small quantities of ammonium hydroxid, 0.02 per cent, greatly retarded the germination of cress. It seems that the active protein of the cell is very sensitive to ammonia.

AMMONIUM HYDROXID

A series of tests was made using from 0.1 to 0.01 per cent of ammonium hydroxid. Four different seeds, cotton, corn, soybeans, and wheat, were allowed to germinate between cloths saturated with the varying concentrations of ammonium hydroxid. It was found that 0.05 or 0.01 per cent proved injurious, while 0.1 per cent prohibited all germination.

Since it was established that ammonia is harmful to seed germination, another test was carried out to study the ammonia produced by micro-organisms. The results of this study are shown in Table X.

TABLE X.—Effect of sugar and of clover on ammonification

Time in 2-day intervals.	Ammonia nitrogen in 100 gm. of soil.		
	No treat- ment.	1 per cent of sugar added.	1 per cent of clover added.
	Mgm.	Mgm.	Mgm.
1.....	1.98	2.0	3.3
2.....		2.1	4.3
3.....		1.96	2.8
4.....		1.4	2.4
5.....		1.4	2.5
6.....	1.90	2.5	2.6
Total.....		11.36	17.9

Since ammonia formation is largely a product of bacterial action, it was thought that sugar or green manure would cause an enormous increase in this substance. The data of Table X show a slight gain in ammonia in the treated soils, but the amount is far too small to affect germination seriously

CARBON DIOXID

It was found that carbon dioxide, when added in large quantities, retards germination but does not cause the seeds to decay. As soon as the carbon dioxide is removed, germination proceeds in a normal manner. In Table XI is given the periodic evolution of carbon dioxide from soil treated with 1 per cent of sugar and 1 per cent of clover.

TABLE XI.—Effect of sugar and clover on carbon-dioxide evolution

Time in days.	Carbon dioxide in 100 gm. of soil.		
	No treatment.	1 per cent of sugar added.	1 per cent of clover added.
	Mgm.	Mgm.	Mgm.
1.....	4.62	22.0	16.02
2.....	6.82	17.2	12.7
3.....	9.46	36.52	22.0
4.....	7.21	37.84	22.75
5.....	7.57	33.97	22.7
6.....	7.74	29.35	24.2
7.....	7.65	26.40	24.42
8.....	9.68	25.30	22.22
Total.....	60.75	228.58	167.01

From the data in this table it is evident that the amount of carbon dioxid evolved in the presence of sugar or clover is far too small to exert a marked effect on germinating seeds.

CALCIUM CARBONATE

It is well known that free acids greatly retard or prohibit germination (3; 4, p. 37). Aside from the direct effect on seeds, an acid reaction may favor the growth of injurious micro-organisms. Accordingly, two series of tests were made, using a neutral and an acid soil with varying amounts of limestone (CaCO_3). The results of the first test are given in Table XII.

TABLE XII.—*Effect of green clover and calcium carbonate on the germination of cottonseed*

No.	Treatment.	Germination.			
		1 week.	2 weeks.	3 weeks.	Relative.
		<i>Per cent.</i>	<i>Per cent.</i>	<i>Per cent.</i>	<i>Per cent.</i>
1	None.....	85	85	85	100
2	1 per cent of clover.....	55	55	55	64
3	1 per cent of clover, 0.1 per cent of calcium carbonate.....	35			40
4	1 per cent of clover, 0.2 per cent of calcium carbonate.....	15			17
5	1 per cent of clover, 0.5 per cent of calcium carbonate.....	15			17
6	1 per cent of clover, 1.0 per cent of calcium carbonate.....	10			11

The data show clearly that limestone in concentrations of from 0.1 to 1 per cent seriously injured the germination of cotton. The seedlings from limed soils died during the first or second week. A second test, similar to the above, was carried out, using acid soil. Here again calcium carbonate seemed to stimulate the injurious factor.

EFFECT OF HEAT

The results of previous tests indicate very strongly the biological nature of the factor injurious to germination. For example, reduced germination is largely associated with the first stages of decomposition. Second, the data seem to exclude the possibility of harmful gaseous products. It is conceivable that in the early stages of decomposition green tissue is favorable to the growth of certain organisms injurious to germination. Accordingly, a series of experiments were made in which the amount and form of green manure applied, the seed, and the biological factors were modified. From 1.5 to 3 per cent of green manure was added. To remove the biological factor, the jars and contents were sterilized in the autoclave at 15 pounds' pressure for two hours. The results of this study were recorded by photographs. Reading from left to right (Pl. LXXXIV, fig. 6), the jars were treated as follows: A, none,

unsterilized; B, 1.5 per cent of green manure, sterilized; C, 1.5 per cent of green manure, unsterilized; D, 3 per cent of green manure, sterilized; E, 3 per cent of green manure, unsterilized. The soil shown in the pots in Plate LXXXIII, figure 5, was treated with green oats, in Plate LXXXIV, figure 6, with green clover. Since the corn and wheat did not show any injury, these illustrations were not reproduced. The data from cotton, clover, and flax are presented in Plate LXXXIV, figures 1, 2, 3, 4, and 5. A glance at the seedlings in the sterilized soil shows conclusively that heat removes or renders inactive the harmful factor. The percentage germination of all crops in the sterilized green-manure soil was equal to that of the untreated controls. Apparently, sterilization has in some way prevented any injury from green-manuring. This is true with 1.5 or 3 per cent of green manure. When repeated, the same results were obtained. These data are given in Table XIII. All of the results point to an injurious agent of biological nature.

TABLE XIII.—*Effect of heat on the germination of cottonseed*

Letter.	Treatment.	Germination.			Relative.
		1 week.	2 weeks.	3 weeks.	
		<i>Per cent.</i>	<i>Per cent.</i>	<i>Per cent.</i>	<i>Per cent.</i>
A	None.....	95	100	100	100
B	Sterilized.....	85	85	85	85
C	1 per cent of clover.....	10	10	10	10
D	1 per cent of clover sterilized.....	80	80	80	80
E	1 per cent of oats.....	35	35	35	35
F	1 per cent of oats sterilized.....	85	90	90	90

SOURCE OF INJURIOUS AGENT

When portions of diseased seedlings are used to inoculate sterilized green-manured soil, the germination of oil seeds is greatly reduced. Numerous tests show that the harmful agent is readily transferred. From the data it must be concluded that the injury to seed germination is biological, probably due to bacteria or fungi. To study the nature of the agent, a series of tests was made with different micro-organisms.

EFFECT OF BACTERIA

In this series of tests bacteria from seed, from green manure, and from soil were studied. From the nature of the seed coat of cotton it is no doubt very rich in a number of bacteria. According to plate counts, the number of micro-organisms on cottonseed is over 122,000 per gram, or an average of nearly 11,000 organisms to one seed. A comparison of the germination of cottonseed free of bacteria and with bacteria, in unsterilized green-manured soil, did not disclose any difference in germination. The bacteria were removed (2) by exposing the seed to the action of hot mercuric chlorid (HgCl_2) or concentrated sulphuric acid (H_2SO_4). The use of sulphuric acid offers an easy and satisfactory method of

removing micro-organisms from cottonseed. The seeds were placed in a large glass-stoppered bottle containing concentrated sulphuric acid and glass beads. After shaking for two minutes, the seeds were removed with a platinum loop and washed in boiled water. From the data it seems that infection is from some source other than the seed.

It has been shown repeatedly that the addition of green manure to soil is followed by an enormous increase in the number of bacteria. Aside from the increase in bacterial food, the green manure carries with it a great number of bacteria (6, 8, 21). Tests with bacteria-free green manures failed to eliminate the injury.

About 16 pure cultures of bacteria were isolated from diseased seeds and green-manured soil. In order to test the effect of these various micro-organisms on germination, sterilized green-manured soil was inoculated with the various species of bacteria and seeded. The tests were carried out in triplicate, using bacteria-free seed of cotton, peanut, and soybeans. Here, again, bacteria failed to show any effect on the germination of oil seeds. In addition to the pure cultures used in the above experiment, a study was made with four laboratory stock cultures, *Bacillus fluorescens liquefaciens*, *B. subtilis*, *B. mesentericus vulgaris*, and *Streptothrix buccalis*. Heavy inoculations of these organisms did not injure the germination of cottonseed or soybeans. This agrees with the results of earlier workers (12, 13, 15, 18)—that is, bacteria grown on rich nitrogenous media do not injure seed germination. An exception to this is noted with cracked or injured seeds.

EFFECT OF FUNGI

From a study of tests carried out with various combinations of sterilized soil, green manure, and seeds free of micro-organisms, it was found that the harmful factor occurs chiefly in soil. The data in Table XIV show very conclusively the position of injury.

TABLE XIV.—Effect of fungi on the germination of cottonseed

No.	Treatment.	Germination.			
		1 week.	2 weeks.	3 weeks.	Relative.
		Per cent.	Per cent.	Per cent.	Per cent.
1	Sterilized soil, 1 per cent of sterilized clover.	20	70	70	100
2	Sterilized soil, 1 per cent of unsterilized clover.....	15	45	45	64
3	Unsterilized soil, 1 per cent of sterilized clover.....				
4	Unsterilized soil, 1 per cent of unsterilized clover.....				

It seems that the harmful agent is found both in soil and in plant tissue, although it is much more prevalent in soil. The results of later tests confirm this statement.

According to many investigators, fungi may injure seed germination (1, p. 30-39; 7, 12, 15). For example, Muth (15) found *Aspergillus niger* harmful to the germination of various seeds, while Atkinson (1, p. 30-39) and Bolley (5, p. 25-27) report a destruction of cotton and flax seedlings by species of *Rhizoctonia* and *Fusarium*.

Since it is established that certain soil fungi are injurious to very young seedlings, the question arises as to the occurrence and growth of parasitic fungi in green-manured soil. An experimental study of the occurrence of fungi in green-manured soil was made. Microscopical examinations of the diseased seeds showed the presence of many fungi on the primary root tip. Although no systematic study was made, some of the forms showed certain characteristics of the genus *Rhizoctonia* and others of the genus *Fusarium*. From portions of the diseased tissue plates were poured. In this way several species of fungi were isolated. These are described under laboratory numbers. All attempts to secure a pure culture of any species of *Rhizoctonia* failed. The various fungi were used to inoculate large tubes and jars of sterilized green-manured soil. The inoculated soil was planted to bacteria-free cottonseed and soybeans. In the soil cultures no injury to germination was noted, except with culture 1. Here from 75 to 100 per cent of the seedlings were killed. Repeated tests with this unknown culture gave similar results. No injury to corn and wheat was noted from inoculations of culture 1, while soybeans and cotton were quickly destroyed.

Since the diseased root tips showed the presence of a *Rhizoctonia*-like fungus, it was arranged to study the effect of certain species of *Rhizoctonia* isolated from other sources. Two strains were employed—one isolated from potatoes, the other from alfalfa. The potato culture was secured from the Department of Plant Pathology of the Wisconsin Experiment Station; the alfalfa culture was supplied by Mr. Fred Jones, of the University of Wisconsin. Table XV gives the results of this test.

TABLE XV.—Effect of *Rhizoctonia* spp. on the germination of cottonseed

No.	Treatment and inoculum.	Germination.			Relative.
		1 week.	2 weeks.	3 weeks.	
1	None, sterilized. Uninoculated.....	Per cent. 75	Per cent. 80	Per cent. 80	Per cent. 100
2	1 per cent clover sterilized. Uninoculated.....	80	85	85	105
3	None, sterilized. Inoculated with <i>Rhizoctonia</i> sp. from alfalfa.....	60	70	70	86
4	1 per cent clover sterilized. Inoculated with <i>Rhizoctonia</i> sp. from alfalfa.....
5	None, sterilized. Inoculated with <i>Rhizoctonia</i> sp. from potato.....	80	80	80	100
6	1 per cent clover sterilized. Inoculated with <i>Rhizoctonia</i> sp. from potato.....	85	85	85	105

Rhizoctonia sp. isolated from alfalfa proved fatal to cotton seedlings. Two weeks after inoculation all of the young plants were dead. On the contrary, a species of *Rhizoctonia* from potato produced no noticeable injury to cotton seedlings. This difference in the action of the two strains of *Rhizoctonia* is very evident from Plate LXXXIII, figure 6, and the data in Table XV. A species of *Rhizoctonia* from alfalfa produced nearly the same effect on soybeans as on cotton, while the germination of corn was not affected.

A study of the optimum conditions for the growth of culture 1 and *Rhizoctonia* sp. from alfalfa showed that about 25° to 30° C. is the most favorable temperature for both of these fungi. The results of a previous study indicate that about 25° C. is the optimum temperature for the growth of the harmful factor. From the data as a whole, it seems very conclusive that the fungus of culture 1 and probably other fungi are the causative agents in the destruction of germinating seeds.

DESCRIPTION OF THE INJURY

Examination of the diseased seeds shows that the injurious factor probably does not attack seeds until after germination. Apparently the fungus attacks the primary root soon after germination. This occurs when the primary root is from $\frac{1}{2}$ to 1 cm. long. The hyphæ pierce the walls of the host, entirely envelop the root, and often penetrate deep within the tissue. In the affected region the tissue loses its form, turns brown in color, and soon rots. Under the microscope these diseased seedling roots are surrounded by a dense mantle of hyphæ, which are often brown-colored.

RELATION OF GREEN MANURE TO INJURY OF OIL SEEDS

Although the evidence at hand does not warrant a definite conclusion, the author suggests the following as a possible explanation for the injury: The green tissue furnishes an excellent medium for the development of fungi. This is especially true in the first stages of decomposition. After one or two weeks in the soil the green manure undergoes certain changes which render it unsuited to the growth of the injurious fungi.

Just why oily seeds should be so sensitive to fungi is not known. It is possible that the oil partly changes to fatty acids in the process of germination (9, 10). According to Schmidt (19, p. 300-303), oil and fatty acids favor the growth of certain fungi. The fungus may produce a fat-splitting enzyme—for example, lipase. This offers a possible explanation for the selective action of the injurious fungi for oil seeds.

SUMMARY

- (1) Green manures may seriously injure the germination of certain seeds.
- (2) This injury is brought about by the action of certain parasitic fungi.
- (3) In the first stages of decomposition of green clover, numerous fungi develop. Some of these fungi are very destructive to seedlings.
- (4) Oil seeds as a class are very easily damaged by fungi. Starchy seeds, on the contrary, are very resistant.
- (5) Cotton seed and soybeans are examples of seeds extremely sensitive to green manuring. The germination of flax, peanuts, hemp, mustard, and clover is reduced in the presence of decomposing plant tissue, but not to as great a degree as that of cottonseed or soybeans. The germination of buckwheat, corn, oats, and wheat is not affected by green manures.
- (6) The damage to oil seeds from green manures is confined largely to the first stages of decomposition. Experimental evidence shows that two weeks after green manure is added it does not cause any serious injury to the germination of oil seeds.
- (7) Small applications of calcium carbonate seemed to increase the injury to germination.
- (8) The rate of germination determines to a certain extent the degree of injury. Slow germination is marked by a high percentage of diseased seedlings.

LITERATURE CITED

- (1) ATKINSON, G. F.
1892. Some diseases of cotton. *Ala. Agr. Exp. Sta. Bul.* 41, 65 p., 25 fig., 1 pl.
- (2) BARRE, H. W., and AULL, W. B.
1914. Hot water treatment for cotton anthracnose. *In Science*, n. s. v. 40, no. 1020, p. 109-110.
- (3) BOKORNY, Thomas.
1912. Einwirkung einiger basischer Stoffe auf Keimpflanzen, Vergleich mit der Wirkung auf Mikroorganismen. *In Centbl. Bakt. [etc.]*, Abt. 2, Bd. 32, No. 20/25, p. 587-605.
- (4) ———
1913. Über den Einfluss verschiedener Substanzen auf die Keimung der Pflanzensamen. Wachstumsförderung durch einige. I. Mitteilung. *In Biochem. Ztschr.*, Bd. 50, Heft 1/2, p. 1-48.
- (5) BOLLEY, H. L.
1901. Flax wilt and flax sick soil. *N. Dak. Agr. Exp. Sta. Bul.* 50, 60 p., illus.
- (6) BURRI, Robert.
1903. Die Bakterienvegetation auf der Oberfläche normal entwickelter Pflanzen. *In Centbl. Bakt. [etc.]*, Abt. 2, Bd. 10, No. 24/25, p. 756-763.

- (7) EDSON, H. A.
1915. Seedling diseases of sugar beets and their relation to root-rot and crown-rot. *In Jour. Agr. Research*, v. 4, no. 2, p. 135-168, pl. 16-26. Literature cited, p. 165-168.
- (8) ESTEN, W. M., and MASON, C. J.
1908. Sources of bacteria in milk. *Conn. Agr. Exp. Sta. Bul.* 51, p. 65-109, fig. 16-23.
- (9) GREEN, J. R.
1890. On the germination of the seed of the castor-oil plant (*Ricinus communis*). *In Proc. Roy. Soc. London*, v. 48, no. 294, p. 370-392.
- (10) ——— and JACKSON, Henry.
1905. Further observations on the germination of the seed of the castor oil plant (*Ricinus communis*). *In Proc. Roy. Soc. London*, s. B, v. 77, no. 514, p. 69-85.
- (11) HENRY, W. A.
1913. Feeds and Feeding . . . ed. 13, 613 p. Madison, Wis.
- (12) HILTNER, Lorenz.
1902. Die Keimungsverhältnisse der Leguminosensamen und ihre Beeinflussung durch Organismenwirkung. *In Arb. Biol. Abt. Land- u. Forstw. K. Gsndtsamt.*, Bd. 3, Heft 1, p. 1-102, 4 fig.
- (13) LAGERVALL, Algot.
1896. Effect of bacteria on germination. (Abstract.) *In Exp. Sta. Rec.*, v. 8, no. 7, p. 566. [1897.] Original article appeared in *Red. Verks. Ultuna Landbrukinst.*, 1895, p. 49-52. 1896. Not seen.
- (14) MIKULOWSKI-POMORSKI, J.
1913. The fertilizing value of the above-ground parts of cereals and leguminous plants. (Abstract.) *In Exp. Sta. Rec.*, v. 31, no. 4, p. 320. 1914. Original article appeared in *Kosmos* [Lemberg], v. 38, p. 929-951. 1913. Not seen.
- (15) MUTH.
1904. Untersuchungen über die Schwankungen bei Keimkraftprüfungen und ihre Ursachen. *In Ber. Landw. Vers. Anst. Augustenb.* 1903, p. 43-48.
- (16) NOBBE, Friedrich.
1876. *Handbuch der Samenkunde* . . . 631 p., illus. Berlin.
- (17) RUSSELL, H. L.
1913. Report of the director, 1911-1912. *Wis. Agr. Exp. Sta. Bul.* 228, 91 p., 37 fig.
- (18) SAJÓ, Karl.
1901. Einige interessante Erscheinungen beim Keimen der Pflanzensamen. *In Prometheus*, Jahrg. 12, No. 587, p. 236-238.
- (19) SCHMIDT, R. H.
1891. Ueber Aufnahme und Verarbeitung von fetten Oelen durch Pflanzen. *In Flora*, Jahrg. 74 (n. R. Jahrg. 49), Heft 3, p. 300-370.
- (20) WEHMER, Carl.
1911. *Die Pflanzenstoffe* . . . 937 p. Jena.
- (21) WIGGER, A.
1914. Untersuchung über die Bakterienflora einiger Kraftfuttermittel in frischem und gärendem Zustande, mit besonderer Berücksichtigung ihrer Einwirkung auf Milch. *In Centbl. Bakt. [etc.]*, Abt. 2, Bd. 41, No. 1/8, p. 1-232. Literatur, p. 228-232.

PLATE LXXXIII

Cotton seedlings, showing the effect of green manures on their growth:

Fig. 1.—*A, B*, Control; *C, D*, 1 per cent of chopped green oats added to the soil; *E, F*, 1 per cent of chopped green clover added to the soil.






Fig. 2.—Effect of planting immediately after plowing under green manure: *A, B*, Control; *C, D*, 0.25 per cent of green manure added to the soil; *E, F*, 0.5 per cent of green manure added to the soil; *G, H*, 1 per cent of green manure added to the soil; *I, J*, 2 per cent of green manure added to the soil.

Fig. 3.—Effect of planting 2 weeks after plowing under green manure. *A, B*, Control; *C, D*, 0.25 per cent of green manure added to the soil; *E, F*, 0.5 per cent of green manure added to the soil; *G, H*, 1 per cent of green manure added to the soil; *I, J*, 2 per cent of green manure added to the soil.

Fig. 4.—Effect of the depth of green manure on germination: *A*, Green manure placed in the bottom of the pot; *B*, green manure placed at the top of the pot; *C*, green manure placed in about the middle of the pot.

Fig. 5.—Effect of sterilized and unsterilized oats used as a green manure: *A*, Control; *B*, 1.5 per cent of oats added and the mixture sterilized; *C*, 1.5 per cent of oats added without sterilization; *D*, 3 per cent of oats added and the mixture sterilized; *E*, 3 per cent of oats added without sterilization.

Fig. 6.—Effect of *Rhizoctonia* sp. on germination in the presence of green manure: *A, B*, Control; *C, D*, sterilized soil treated with green manure; *E, F*, sterilized soil inoculated with *Rhizoctonia* sp. from potatoes; *G, H*, sterilized soil treated with green manure and inoculated with *Rhizoctonia* sp. from alfalfa.









PLATE LXXXIV

Clover, flax, and cotton seedlings, showing the relation of green manures to germination in sterilized and unsterilized soil:

Fig. 1.—Clover: *A*, control; *B*, 1.5 per cent of chopped green oats added and the mixture sterilized; *C*, 1.5 per cent of chopped green oats added and the mixture not sterilized; *D*, 3 per cent of chopped oats added and the mixture sterilized; *E*, 3 per cent of chopped oats added and the mixture not sterilized.

Fig. 2.—Clover: *A*, control; *B*, 1.5 per cent of chopped clover added to the soil and the mixture sterilized; *C*, 1.5 per cent of chopped clover added to the soil and the mixture not sterilized; *D*, 3 per cent of chopped clover added to the soil and the mixture sterilized; *E*, 3 per cent of chopped clover added to the soil and the mixture not sterilized.

Fig. 3.—Flax: *A*, control; *B*, 1.5 per cent of chopped oats added to the soil and the mixture sterilized; *C*, 1.5 per cent of chopped oats added to the soil and the mixture not sterilized; *D*, 3 per cent of chopped oats added to the soil and the mixture sterilized; *E*, 3 per cent of chopped oats added and the mixture not sterilized.

Fig. 4.—Flax: *A*, control; *B*, 1.5 per cent of chopped clover added and the mixture sterilized; *C*, 1.5 per cent of chopped clover added to the soil and the mixture not sterilized; *D*, 3 per cent of chopped clover added to the soil and the mixture sterilized; *E*, 3 per cent of chopped clover added to the soil and the mixture not sterilized.

Fig. 5.—Cotton: *A*, control; *B*, soil sterilized; *C*, 1 per cent of chopped clover added to the soil and the mixture not sterilized; *D*, 1 per cent of chopped oats added to the soil and the mixture not sterilized; *E*, 1 per cent of chopped clover added to the soil and the mixture sterilized; *F*, 1 per cent of chopped oats added to the soil and the mixture sterilized.

Fig. 6.—Cotton: *A*, control; *B*, 1.5 per cent of chopped clover added to the soil and the mixture sterilized; *C*, 1.5 per cent of chopped clover added to the soil and the mixture not sterilized; *D*, 3 per cent of chopped clover added to the soil and the mixture sterilized; *E*, 3 per cent of chopped clover added to the soil and the mixture not sterilized.

A NEW SPRAY NOZZLE

By C. W. WOODWORTH,

Entomologist, Agricultural Experiment Station of the University of California

INTRODUCTION

A new principle has been discovered in nozzle construction whereby a flat spray can be produced with a uniform distribution of the water comparable to that of the hollow cone of spray from a cyclone nozzle. Hitherto all flat sprays have been of lenticular section, breaking up into fine mist on the sides and into relatively coarse drops in the center. It was observed that the flat spray produced by two impinging streams was at right angles to the original plane of motion of the two streams, but when the streams failed to meet squarely the plane was shifted and could, in fact, be moved through an arc of 180° with a very great change in the distribution of the water currents. It requires only a slight angular deviation to decrease very perceptibly the coarseness of the central drops, producing greater uniformity, and a position can be reached in which the coarsest drops are on the edge, those in the center therefore being the finest.

The principle finally discovered was that when two streams meet across half their section the resulting sheet of spray will be of practically uniform thickness throughout, occupying a plane 45° from the plane of the streams and finally breaking up into drops of great fineness and uniformity.

PRODUCTION OF SPRAY

There are two causes that may act in the production of spray particles: (1) Friction, which may cause an eddy along the edge of the stream sufficient to break the surface tension and allow the small eddying masses to fly off from the column of water; and (2) divergence of the direction of motion of the particles, resulting in the thinning out of the water mass in the form of irregular sheets until the surface film finally gives way and the sheet of water is suddenly converted into drops.

Both methods may be seen in the breaking up of the stream from a simple nozzle where, from the sides of the solid column of water, very minute particles of mist are given off, while the velocity and friction are great. With decreasing velocity farther on the eddies become larger, the mist gradually becomes coarser, and, finally, as the spread of the stream makes it break up into irregular sheets of water, the size of the drops produced by the second process results in an intermingling of drops of all sizes. At first the drops are very accurately graduated,

those of the same size being produced at the same distance from the nozzle, but when the second process replaces friction as a cause of spray production, irregularity results, owing to the irregular shapes of the water sheets.

In a cyclone nozzle the stream at once diverges widely in the form of a hollow cone. Friction plays no part in the production of the spray, but the cone increases so rapidly in diameter that the liquid soon becomes a very thin sheet of unvarying thinness all the way around, and breaks into a uniformly fine mist. The uniformity may be assumed from the fact that on all sides the sheet extends an equal distance from the orifice before breaking into a spray, and experimentally can be shown to exhibit to an equally high degree both fineness and uniformity.

Figure 1 expresses in a diagrammatic form the facts shown by the photographs. The circles show the actual positions of the orifices in




FIG. 1.—Diagram showing the characteristic differences between the three forms of impinging-stream nozzles.

each case and the black transverse marks give the effect of the impinging streams; the water remains thickest in the middle in C, thickest at the edges in A, while in B it is spread out evenly.

Above, the black portion indicates the water sheet, the sizes of the spots along the margin indicate the sizes of drops produced at these points, and the approximate velocity of the drops is shown by the length of the lines radiating from these spots.

SPRAYS PRODUCED BY IMPINGING STREAMS

The actual movement of the water in forming a spray through the impact of two streams is shown in Plate LXXXV and Plate LXXXVI, figure 1. It was not found practicable to secure the successive pictures with sufficient rapidity to show more than two steps in the forming spray, but by interpolating, a fairly satisfactory series was obtained. The

right-hand nozzle is of the common type where the streams impinge squarely. The middle nozzle is of the new type, but not strictly comparable with the former, since the streams come together at a broader angle, making a wider spray. Indeed, when the spray is under full pressure (Pl. LXXXVI, fig. 1) the spread is too wide, producing a lateral dribble and marginal fringe of spray. The left-hand spray is intermediate in angle and spread and gives the fish-tail effect.

The contrast is shown from the first illustration, the fish tail having thick marginal zones and the other two thick central zones, much shorter in the middle nozzle. In Plate LXXXVI, figure 1, where the spray sheets assume their normal proportions under high pressure, the large size of the white patch in the middle corresponds to the better final distribution of the spray particles. The irregularity of the spot shown on the left of this white patch is due to an irregularity in the orifice on the opposite side.

In Plate LXXXVI, figure 2, which shows the result of a sudden decrease of pressure, the character of the water sheets becomes especially evident, since they are increased greatly in size and the production of spray almost ceases.

ADVANTAGE OF A FLAT SPRAY

The cyclone nozzle leaves nothing to be desired in the way of fineness and uniformity of spray, but it has the disadvantage of making a ring of spray which surrounds instead of touching the object towards which the nozzle is directed. It is very difficult for one handling the nozzle to keep in mind the fact that the spray is strictly limited to the visible parts of the cone. A flat spray, on the other hand, reaches the point aimed at and is more available for treating branches of trees, for example, where the desire is to concentrate the spray on a line. For general spraying also the use of a flat spray, like the use of a flat brush for painting, gives uniform results more quickly and easily. For these reasons, while no other nozzle on the market produces a flat spray comparable in quality to the spray produced by the various types of cyclone nozzles, they are, nevertheless, more extensively used than the cyclone nozzles.

ADVANTAGE OF UNIFORMITY AND FINENESS

The use of nozzles of the flat type is generally acknowledged to be for the purpose of securing the flat shape of spray fan and is not a rejection of the principle that a uniformly fine spray is the most desirable. In fact, the use of these nozzles is generally associated with the use of high pressures, whereby the defects of a poor grade of nozzle are less apparent. The particular advantage of fineness is that it makes possible the even distribution of the spray material.

Fineness involves evenness. In a nozzle giving coarse drops, part of the material is in a finely divided state, and the improvement in a spray

nozzle comes through decreasing the size of all but the smallest particles and thus increasing the proportion of minute particles until, as in the cyclone nozzle, practically all of the material is in the most finely divided state and is therefore also uniform. This improvement can be produced by increasing the pressure or decreasing the size of the stream. Under the same pressure a nozzle with a large orifice gives coarser drops than a similar nozzle with a small orifice. Therefore, where a larger volume of spray is desired, it has been the practice to duplicate the nozzles rather than enlarge them, giving clusters of nozzles; but where high pressure is available, large nozzles, particularly those of the better type, may be used. With extreme pressures, such as were employed in the gipsy-moth work and in the walnut spraying in California, a nozzle of the poorest quality and rather large size has proved to be practical. In nearly all cases the desirability of fine and uniform sprays, in order to secure evenness of distribution, has been recognized. It is possible, however, that under some circumstances a driving spray may be desirable, and this can be secured only by the use of less efficient nozzles.

VARIATION IN FINENESS

The sizes of the smallest drops in a spray are not necessarily the same, particularly when made by the breaking up of a sheet of water. By a change in the proportions of the eddy chamber in a cyclone nozzle or by a change in spraying pressure the diameter of the cone at the point of breaking can be changed, and the drops will remain uniform, but will be of a different size than before. In the new type of nozzle here described the angle of impact and the spraying pressure exert similar effects, and a series of nozzles can be produced covering much the same range obtainable in a cyclone nozzle and distinguishable by the width and length of the fan.

Only relatively small drops in the spray in either case are obtained, and these show great uniformity, the variation in size being inside of rather narrow limits.

The new type of nozzle is the form in which the spray is in a plane inclined at the angle of 45° from the plane of the impinging streams, but between that and the usual style, having the spray in a plane 90° from that of the streams, there is the possibility of any number of intermediate forms that present any desired degree of uniformity in the size of the drops. Should a compromise nozzle giving a driving spray with greater uniformity than in the existing nozzles be desired, it can readily be constructed. The same could be secured by a disproportion between the sizes of the two streams, and in this case the coarser portion would be at one edge instead of at the center of the fan. This form might be desirable for some spot-spraying for scale insects, and it might be desirable to have a means of controlling the size of one of the streams.

WHERE THE NEW NOZZLES ARE IMPRACTICAL

Because the spray must first be separated into two streams in this type of nozzle it becomes particularly liable to clogging and should not be used for any spraying where there is any such tendency—e. g., with Bordeaux mixture.

Most of the spray materials now used, however, are clear solutions and give no trouble in the nozzle.

LONG- AND SHORT-DISTANCE NOZZLES

When the angle is widest between the impinging streams, the angle of the fan is likewise widest, the drops finest, and the carrying distance of the spray the shortest.

An acute angle between the impinging streams produces a very narrow spray which carries a longer distance, but may perhaps finally reach nearly as great a width as that of the rapidly spreading short-distance spray.

Some prefer a long-distance nozzle and use it close to an object, as where spot spraying on a tree trunk is desired. The new type of nozzle lends itself very readily to adjustment to any degree of distance, from the shortest to nearly the longest found in spray nozzles.

ADJUSTMENT

Any form of two-stream nozzle, like that known as the calla, or lily, nozzles, can be quickly converted into a nozzle of the new type by the use of a reamer, slightly enlarging the two apertures on opposite sides by working the instrument obliquely to the surface of the nozzle and trying it from time to time until the spray sheet stands at 45° .

The same process will enable one to adjust a nozzle at any time should it wear irregularly enough to change the angle of the spray fan. The shape of the fan is a good index of the correct adjustment. If the angle is just right, the fan is triangular; if less than 45° , it is shortest in the center and the spray is coarser at the ends. If the angle is more than 45° , the fan is longest in the center and the spray coarsest at this point.

With care the reamer can be so used as to effect the change in the stream without enlarging the hole at the surface, and, therefore, not changing the volume of discharge. It may be possible to change the angle of the spread of the fan by reaming out beneath on the side adjacent to or opposite the other hole. One should continually try a nozzle while adjusting it, so as not to carry the work too far.


SUMMARY

- (1) A new principle employed in nozzle construction will produce a flat spray with the qualities of a cyclone nozzle.
- (2) A uniform sheet of water breaking along its edge produces drops of uniform size.

-
- (3) A flat spray is more easily directed and produces a more uniform distribution than the cone of spray from a cyclone nozzle.
 - (4) Uniformly fine drops of spray aid in securing uniformity of distribution.
 - (5) The new nozzle allows some variation in size of spray.
 - (6) It also may be made into a long- or short-distance nozzle.
 - (7) It can be easily constructed by modifying existing nozzles and may be adjusted if it becomes worn.

PLATE LXXXV

The beginning of the spray from three kinds of nozzles, as photographed with a moving-picture camera.






PLATE LXXXVI

Fig. 1.—The appearance of spray from three kinds of nozzles as full pressure is applied (a continuation of Plate LXXXV).

Fig. 2.—Two stages at the end of the spray as the pressure is reduced.

A NEW INTERPRETATION OF THE RELATIONSHIPS OF TEMPERATURE AND HUMIDITY TO INSECT DEVELOPMENT

By W. DWIGHT PIERCE,

Entomological Assistant, Southern Field Crop Insect Investigations, Bureau of Entomology

INTRODUCTION

Upon the proper interpretation of the laws of climatic control of life rests the solution of many practical problems, and inasmuch as all plant and animal life reacts to climate in the same general manner it is apparent that the study of the climatic control of insect development may throw light upon the problems of all other forms of life. It has been apparent to some workers in the field of ecology that our so-called laws of effective temperature were deficient in many respects. A large number of phenomena were not properly explained by any known theory. It is with the hope that the present interpretation may come closer to the truth that this paper has been prepared.

Biologists for years were laboring with the theory of a fixed zero of effective temperature for all life, and only recently was it accepted that each species might have a different zero. It has been the custom to determine the thermal constant for any given activity by multiplying the number of effective degrees accumulated above the effective zero in daily units of mean temperature by the time in which the given phenomenon took place. The noneffective low temperatures were eliminated, but not the time in which they were experienced. Inasmuch as most workers were located in north-temperate climates, where high noneffective temperatures seldom occur, it had not occurred to them that some high temperatures might not be effective and that there was another boundary to the effective zone besides the zero. These high temperatures and the time in which they are experienced must be eliminated. In addition to all of these errors in method, there has been no correlation of the humidity factor until very recently, although now many workers are trying to solve the part played by this factor.

The principal data upon which the writer has based his studies include records of thousands of individual boll weevils (*Anthonomus grandis* Boh. and *A. g. thurberiae* Pierce), made by the members of the boll-weevil force under the direction of Mr. W. D. Hunter and the writer at various localities in Texas, Louisiana, and Arizona throughout the period of years from 1902 to 1915. At each place where biological notes were made a thermograph-hygrograph record was kept, and this record was

checked twice daily by maximum and minimum thermometer and sling-psychrometer readings. The means of temperature and humidity are based upon these records. In addition to the natural records, a series of artificial-cold experiments were conducted at various times, and the writer recently conducted an extensive series of artificial-heat experiments with definite humidity control in order to determine the effects of heat.

EXPERIMENTAL METHODS

Before venturing to present this new interpretation the writer has thoroughly discussed it with many prominent workers, and it is now proposed for more extensive criticism and elaboration.

To express the relationship of the two factors, temperature and humidity, to insect metabolism, development, and activity, a temperature scale may be marked off on the vertical line of a sheet of plotting paper and a humidity scale from left to right on the horizontal line. There are, for any given insect, definite boundaries of atmospheric temperature and humidity within which the life of the species revolves. There is a temperature below which, even for the shortest time, life is impossible—the absolute minimum fatal temperature. There is also a temperature above which, even for a moment, life is impossible—the absolute maximum fatal temperature. Absolute dryness is more or less prohibitive of life and so is absolute humidity, or saturation, although some insects may be adapted better to withstand extremes of humidity than others. It is quite possible that the boundaries of humidity may be 0 and 100 per cent, or infinitesimally close thereto.

The diagrammatic figure sought, however, has four definite absolute boundaries—the maximum and minimum temperatures and humidities.

Within the limits which we have thus defined there exist conditions under which all the activities of the species reach their maximum efficiency. It has been conceived by most writers that this maximum efficiency was reached at a definite point known as the optimum. It seems more likely that it will prove to be a zone of humidities and temperatures of more or less restricted area. A careful study of the records of any species, charting for the time required for each activity and the temperature and then similarly for humidity, will disclose temperature and humidity points of maximum efficiency. With the boll weevil these points lie approximately near 83° F. and 65 per cent of relative humidity.

ZONES OF CLIMATIC RELATIONS

At any ordinary humidity, starting with the absolute minimum fatal temperature, as the temperature increases a longer and longer time of exposure is required to kill, until a point is reached at which life continues indefinitely. This zone of temperatures has been called the zone of fatal temperatures.

As the temperature continues to rise it passes through a zone of ineffective temperatures, known commonly as the zone of hibernation, which the writer will shortly prove to be an inappropriate term. At the lowest temperatures in this zone complete dormancy without metabolism is found; but as the temperature increases a gradual approach to sensibility is noted, first metabolism, next movement, and then the necessity of feeding. The point at which metabolism or growth begins at a given humidity is the zero of effective temperature.

As the temperature increases above this zero the activity is at first very sluggish, but becomes more and more active until the so-called optimum is reached, and from this point upward the temperatures cause less and less activity, inducing stupor and finally sleep or coma.

At the point of coma begins the zone of ineffective temperatures formerly known as estivation. With the increase of temperature sleep becomes more and more sound until a point is reached at which death occurs after long exposure. At this point begins the zone of high fatal temperatures at which death occurs at shorter and shorter periods until it is instantaneous at the absolute maximum fatal temperature. This completes the vertical cross section of the figure desired. A statement regarding these vertical zones was first published by the Bureau of Entomology in 1912.¹

In the past, however, the fact that a similar horizontal cross section at any temperature can be made, starting at absolute dryness and reading toward absolute humidity, has not been recognized. In this manner are shown zones of fatal dryness, dryness causing stupor, increasingly effective humidity, the most effective humidity, decreasingly effective humidity, excessive humidity causing drowsiness, and finally fatal humidity, at least under certain conditions of exposure.

In the case of the boll weevil the resulting figure is a series of concentric ellipses centered about the optimum and with diagonal axes. On the accompanying diagram the main details of the relations of temperature and humidity to the boll weevil are brought out. Only a few of the more salient records are included. The development in buds (cotton squares) is based upon hundreds of individual records, but is not reported in detail. The outer lines are much less definitely located than the inner ones, but whatever their actual location the picture would be substantially the same.

EFFECTIVE TEMPERATURE

Workers who have used the zero of effective temperature in their studies will note that, according to the present theory, the zero when charted is an elliptical curve representing a different point at each degree

¹ Hunter, W. D., and Pierce, W. D. Mexican cotton-boll weevil. 64d Cong., 2d Sess., Sen. Doc. 305 (U. S. Dept. Agr. Bur. Ent. Bul. 114), p. 125-128. 1912.

of humidity. Because of the difficulty of computing this zero, the writer has been requested to describe his method of computing effective temperatures.

The first step is to tabulate all records of a given mean percentage of humidity on a single sheet. The zone of effective temperatures must be worked out separately at each degree of humidity. Only by a laborious series of testings can the first zero be approximated, unless the worker finds it by a fortunate chance. The total effective temperature is the criterion by which we finally know when we have rightly defined the limits of the zero. This is known as the thermal constant and is the multiple of the mean of the effective temperatures (between the zero and the absolute), figured in day units, by the time in which these effective temperatures were experienced. Noneffective temperatures, whether high or low, and the time in which they were experienced must be eliminated. The zone of effective temperatures will be finally reached for any given humidity when the




FIG. 2.—Graph showing the method of determining the zone of effective temperatures at a humidity of 50 per cent.

temperatures will be finally reached for any given humidity when the

difference in the total effective temperatures is reduced to a minimum. At the start some arbitrary zero must be chosen and the effective temperatures computed above this. Then it is necessary to remove degree by degree at the top or bottom and note each time whether the difference in the total effective temperatures becomes larger or smaller. This process may be charted so that the general tendency can be seen. The figures found in the writer's attempt to establish the zone of effective temperatures for the boll weevil at 56 per cent humidity will illustrate the manner in which the points desired were ascertained. These results are presented in figure 2, and it will be seen that the first tentative zone chosen was 51° to 100° F. By much testing it was narrowed to within the limits of 75° to 92° F., for which the optimum is practically 83.5°.

Having obtained the limits of the zone, the records of development in cotton squares at a mean humidity of 55.9 per cent to 56.9 per cent, made at Victoria, Tex., in 1913, by Mr. B. R. Coad, of the Bureau of Entomology, are as shown in Tables I and II.

TABLE I.—Records of development of *Anthonomus grandis* at Victoria, Tex., in 1913, at a humidity of 55.9 to 56.9 per cent

Experiment.	Mean humidity.	Date of oviposition.	Time of maturing.	Actual period of development.	Number of weevils observed.		Total weevil days.	Actual temperature.		
					Male.	Female.		Absolute maximum.	Absolute minimum.	Mean.
	Per cent.			Days.				° F.	° F.	° F.
1.....	56.1	July 27	Aug. 9	13	6	2	104	104	73.2	88.2
2.....	56.4	July 26	Aug. 8	13	1	3	52	104	73.2	88.2
3.....	56.6	July 27	Aug. 10	14		1	14	104	73.2	88.2
4.....	56.9	July 27	Aug. 11	15		1	15	104	73.2	88.2
5.....	55.9	May 22	June 7	16	1		16	95.5	54.5	78.2
Mean.....	56.2				Total 8	7	201			

TABLE II.—Records of development of *Anthonomus grandis* at Victoria, Tex., in 1913, in the zone of effective temperatures, 75° to 92° F.

1a	2a	3a	4a	5a	6a	7a	8a	9a	10a	11a
Experiment.	Number of weevils.	Mean humidity.	Humid time units.	Period experiencing effective temperature.	Total effective weevil days.	Mean effective temperature.	Effective thermal units.	Mean daily effective temperature, units.	Total effective temperature.	Humidity plus effective temperature.
		Per cent.		Days.		° F.			° F.	° F.
1.....	8	56.1	448.8	8.19	65.52	83.8	670.4	8.8	72.0	139.9
2.....	4	56.4	225.6	8.15	32.72	83.8	335.2	8.8	71.98	140.2
3.....	1	56.6	56.6	8.86	8.86	83.6	83.6	8.6	76.1	140.2
4.....	1	56.9	56.9	9.52	9.52	83.7	83.7	8.7	82.82	145.5
5.....	1	55.9	55.9	9.95	9.95	82.0	82.0	7.6	75.6	118.5
Total.....	15		843.8		126.57		1,255.5	8.7	73.5	139.9
Average.....		56.2			8.43		83.7			
Difference.....								1.2	10.84	2.1

a Column 4 is product of columns 2 and 3. Column 5 is computed from the actual records. Column 6 is the product of 2 and 5. Column 8 is the product of 5 and 7. Column 9 is 7 minus the zero (75° F.). Column 10 is the product of columns 5 and 9. Column 11 is the sum of columns 3 and 7.

From these tables it will be seen that the effective period of development is from 8 to 10 days, averaging 8.43 days, while the actual development ranged from 13 to 16 days. It is noticeable that in all of the records the maximum as well as the minimum temperatures ran outside of the zone of effective temperatures. The total effective temperature ranged from 72° to 83° F., with 73.3° as the weighted mean and with a total difference of only 10.84°, a very small difference.

It is not necessary in this paper to give the further details of the zone of effective temperatures at other humidities. The determination of the zone for the next percentage of humidity is much less difficult, because it must be just a little narrower or a little wider than already determined. As the axis is diagonal, the upper and lower bounds will depart at a different rate. After several points have been determined, the axis can be located and then the figuring becomes very simple. It must be noted that every hour of effective temperature has its cumulative effect, even in the winter time.

ZONE OF INACTIVITY

One of the results of the acceptance of the present interpretation will be the necessity of discarding the conception of separate zones of hibernation and estivation. Physiologists have demonstrated that the effects of heat and cold on metabolism are alike. The writer has frequently noticed in field work the impossibility of differentiating between a frozen and a heat-killed boll-weevil larva. Prof. G. G. Becker, of Arkansas Agricultural College, several years ago observed that the fall army worm, *Laphygma frugiperda* S. and A., had two periods of activity and two of inactivity every day in the hot days in the Ozarks. Activity began in the morning and continued until the early part of the afternoon, when the heat caused the worms to be inactive for several hours. They then again became active during the early hours of the night, but the nights were cold and the worms became inactive until morning. The phenomena of a year were reproduced day by day. Inactivity due to cold in the summer time can not properly be called hibernation.

In Arizona the boll weevil is now native on wild cotton (*Thurberia thespesioides*). It normally breeds in the bolls in the fall, becoming adult by December 1, but remains in its cell throughout the cold winter and the warming spring. In some canyons there is a spring rainy season and *T. thespesioides* has a spring fruiting season. In these localities the moisture also releases the weevils from their cells and they begin breeding. A dry season follows and the weevils go to sleep. In other canyons the spring is not wet and the plants and weevils are inactive until the regular rainy season in August, when the long rest is broken. In some canyons the weevils therefore have two resting periods during the year, and in other canyons they are at rest from fall until summer. It not infrequently happens that the August rainy season does not materialize, and under

such circumstances the weevils stay in their cells and the plants remain dormant until the next year or perhaps for several years. As evidence of this the writer kept several of these weevils over 500 days without food or water, and one lived 626 days, dying only when moisture invaded the room where it was kept.

Hunter, Pratt, and Mitchell¹ record the unusual ability of larvæ of *Hermetia chrysopila* Loew, a cactus scavenger fly, to withstand long periods of drought. Larvæ in various stages of development were kept for more than 15 months without food and developed readily later when food was supplied. The very leathery integument seems to protect the insect against desiccation, and in other ways the larva has evidently adapted itself to long periods of waiting for favorable food, which, in the arid regions, depends upon the infrequent rains. Both of these instances are more properly resting periods due to dryness than to cold or heat.

NOMENCLATURE OF CLIMATIC EFFECTS ON LIFE

As charted, there are three elliptical zones which express the three principal effects of climate on life, viz, activity, inactivity, and death. The zone of activity may be known as the "thermopractic" zone (θερμός, meaning heat, plus πρακτικός, meaning effective). The zone of inactivity may be known as the zone of "anesthesia" (άναισθησία, meaning insensibility). The zone of death may be known as the "olethric" zone (ολέθριος, meaning deadly). The region of greatest activity may be known as the "practicotatum" zone (πρακτικώτατον, meaning most effective).

Many phases of climatic effects have been differentiated, and medical literature abounds in words descriptive of these effects. For some effects no words are available. The writer has thought it best to present a complete and consistent system of nomenclature, based on the Greek, using all words already in the language, and only supplying new words where none are now available.²

It may be convenient to refer to the most effective temperature or the most effective humidity, in which cases we may use the words "thermopracticotatum" or "hygropracticotatum."

The awakening from sleep is termed "anastasis" (ανάστασις). We can therefore speak of "thermanastasis" and "hygranastasis," depending on whether the awakening is caused by a change of temperature or a change of humidity.

Heat, moisture, dryness, or cold added to the "practicotatum" will cause sluggishness. We have to indicate this condition the term "nochelia"

¹ Hunter, W. D., Pratt, F. C., and Mitchell, J. D. The principal cactus insects of the United States. U. S. Dept. Agr. Bur. Ent. Bul. 113, p. 38-39. 1912.

² New Standard Dictionary. 1913.

Conk, C. M. An Illustrated Dictionary of Medicine, Biology and Allied Sciences . . . ed. 6, with . . . Sep . . . 1623, 571 p., Philadelphia, 1910.

(*παχέλεια*, meaning sluggishness) and can show the type of sluggishness by the addition of a prefix, as "thermonochelia," "hygronochelia," "xeronochelia," and "rhigonochelia."

At least three of these factors produce under extreme conditions a stifling sensation, and we may express this by the terms "thermopnigia," "xeropnigia," and "hygropnigia" (*πνίγος*, meaning stifling).

The stifling sensation ends in complete insensibility, or anesthesia, and this word may be modified to express the cause, as in the term "thermanesthesia," "hygranesthesia," "xeranesthesia," and "rhiganesthesia."

Death from heat is known as thermoplegia (*πληγή*, meaning stroke), while from excessive moisture it may be known as "hygroplegia," and from freezing, as "rhigoplegia." Death from drying is known as "apoxeraenosis" (*ἀποξηραίνω*, meaning to dry up).

The determination of locomotion by heat is called "thermotaxis," and movement brought about by heat is called "thermotropism."

Unusual sensibility to heat is called "thermalgesia" and "hyperthermalgesia." The ability to recognize changes of temperature is "thermesthesia," and its extreme is designated as "thermohyperesthesia," abnormal sensitiveness to heat "stimuli." Fondness for heat or requiring great heat for growth is called "thermophilic," while resistance to heat is called "thermophylic." Rapid breathing, owing to high temperature, is designated as "thermopolypnea," contraction under the action of heat as "thermosystaltic," adapting the bodily temperature to that of the environment as "pecilothermal," and a morbid dread of heat as "thermophobia."

The life after apparent death, called "anabiosis," is exemplified in such cases as that of the *Hermetia* larvæ mentioned above.

Pain from the application of cold is called "cryalgnesia," abnormal sensitiveness to cold "cryesthesia," and a morbid sensitiveness to cold "hypercryalgnesia."

PRACTICAL APPLICATIONS

Many practical measures will result from the further study of climatic relations to life. A few of these may be indicated.

One of the most effective measures for the control of the cattle tick is pasture rotation based upon the possible duration of life of the seed tick without an animal host. As this period varies with the season, it is necessary to know the climatic laws under which this species reacts.

The fall army worm advances across the country and again retreats in complete accord with changing temperatures. The proper fixation of the zone of effective temperature may make it possible to plan the planting of winter crops to avoid damage.

The cotton boll weevil must have food up to the time that it enters hibernation. Early harvesting and destruction of stalks before the low temperatures set in offer one of the most satisfactory methods of control.

ADDITIONAL COPIES
OF THIS PUBLICATION MAY BE PROCURED FROM
THE SUPERINTENDENT OF DOCUMENTS
GOVERNMENT PRINTING OFFICE
WASHINGTON, D. C.
AT
15 CENTS PER COPY
SUBSCRIPTION PRICE, \$3 PER YEAR

Δ

INDEX

	Page		Page
<i>Acer</i> —		Announcement of Weekly Publication.....	i
<i>rubrum</i> , host plant of <i>Comandra umbellata</i> ...	134	<i>Antennaria plantaginifolia</i> , host plant of <i>Comandra umbellata</i>	134
<i>saccharinum</i> , composition of sap of.....	538-540	<i>Anthonomus grandis</i> —	
<i>saccharum</i> —		development of.....	1188-1189
composition of sap of.....	538-540	relation of—	
mineral composition of.....	540-543	humidity to development of.....	1183-1191
<i>Achillea millefolium</i> , host plant of <i>Comandra umbellata</i>	134	temperature to development of.....	1183-1191
Acid, fruit, toxicity of, to <i>Sclerotinia cinerea</i> ..	388	<i>Apanteles militaris</i> —	
<i>Acinomyces</i> —		biology of.....	495-508
<i>horis</i> , effect of low temperature on.....	654	endoparasite of <i>Heliothrips unipunctata</i>	495
<i>chromogenus</i> , effect of low temperature on.....	651-652, 654	function of caudal vesicle of.....	504-506
<i>organicus</i> , effect of low temperature on.....	651-652	life stages of.....	495-501
Activity of Soil Protozoa (paper).....	477-488	origin of caudal vesicle of.....	504-506
Aeration, effect of, on germination.....	1168	Aphis, green apple. See <i>Aphis pomi</i> .	
Age, relation of, to occurrence of tumors in domestic fowl.....	399	<i>Aphis</i> —	
Agglutination Test as a Means of Studying the Presence of Bacterium abortus in Milk (paper).....	871-875	<i>pomi</i> —	
Air, agency in dissemination of <i>Cercospora personata</i>	895-897	biology of.....	955-994
Alcohol, combustion of, in respiration calorimeter.....	345	diomorphic reproduction by.....	918
Alfalfa. See <i>Medicago sativa</i> .		distribution of.....	957-958
Alkali salts—		egg stage of.....	960-967
combinations of, effect on plant growth.....	13-48	feeding habits of.....	981-984
effect of, on germination and growth of crops.....	1-53	forms of.....	980-981
toxicity of.....	1-53	history of.....	957-958
Allard, H. A. (paper), Distribution of the Virus of the Mosaic Disease in Capsules, Filaments, Anthers, and Pistils of Affected Tobacco Plants.....	251-256	life of stem mother of.....	968-970
<i>Alternaria</i> —		methods of studying.....	958-960
<i>panax</i> , parasite of <i>Panax quinquefolium</i> ..	181-182	morphology of.....	955-994
<i>solanii</i> , effect of low temperature on.....	652	nomenclature of.....	956
<i>Alternaria panax</i> , the Cause of a Rootrot of Ginseng (paper).....	181-182	overlapping generations of.....	981-982
Alway, F. J., and Bishop E. S. (paper), Nitrogen Content of the Humus of Arid Soils.....	900-916	sexes—	
<i>Amaranthus retroflexus</i> , transpiration of.....	618-623	habits of.....	984-991
<i>Ammonium hydroxid</i> , effect of, on germination.....	1168-1169	life of.....	984-991
<i>Anerobe</i> —		summer forms of.....	970-978
<i>facultative</i> —		<i>spp.</i> , plan of description of.....	967-968
non-spore-forming.....	939	Apparatus for Measuring the Wear of Concrete Roads (paper).....	951-954
slime-forming.....	940	Apple. See <i>Malus</i> .	
<i>Andropogon</i> —		<i>Arachis hypogaea</i> —	
<i>sorghum</i> , transpiration of.....	597-602	composition of seed of.....	1162
<i>virginicus</i> , host plant of <i>Comandra umbellata</i>	134	lealspot of.....	891-902
<i>Angelica silosa</i> , host plant of <i>Comandra umbellata</i>	134	Army worm, fall. See <i>Laphygma frugiperda</i> .	
Angular Leafspot of Cucumbers (paper)....	465-476	Arsenic—	
Anions, effect of, on growth of <i>Triticum</i> spp.	42-43	absorption of, by soil.....	460
		applied as spray for weeds, late and effect of.....	459-463
		fixation reactions in soil.....	461-463
		in soil, effect of irrigation on.....	460-461
		<i>Arrhenatherum elatius</i> , syn. <i>Avena elatior</i> .	
		<i>Ascochyta colorata</i> , effect of low temperature on.....	654
		Ash Composition of Upland Rice at Various Stages of Growth (paper).....	357-364
		Asiatic Species of Gymnosporangium Established in Oregon, An (paper).....	1003-1010

	Page		Page
Asphalt—		Baker, A. C. (paper), Identity of <i>Eriosoma</i>	
cement. See Cement, asphalt.		pyrii.....	1115-1119
effect of controllable variables upon the		Baker, A. C., and Turner, W. F. (paper),	
penetration test for.....	805-818	Morphology and Biology of the Green	
oil, results of needle standardization test...	1123	Apple Aphid.....	955-994
Aster—		Balance—	
<i>ericoides</i> , host plant of <i>Comandra umbellata</i>	134	automatic recording step-by-step.....	117-120
<i>macrophyllus</i> , host plant of <i>Comandra</i>		continuous-record.....	120-122
<i>umbellata</i>	134	Ballinger, A. M., and Johnson, P. M. (paper)	
<i>patens</i> , host plant of <i>Comandra umbellata</i>	134	Life-History Studies of the Colorado Potato	
<i>undulatus</i> , host plant of <i>Comandra umbel-</i>		Beetle.....	917-926
<i>lata</i>	134	Banana as a Host Fruit of the Mediterranean	
<i>Atriplex hortensis</i> , host plant of <i>Peronospora</i>		Fruit Fly (paper).....	793-804
<i>effusa</i>	59-67	Banana. See also <i>Musa</i> spp.	
Aubel, C. E., and Wentworth, E. N. (paper),		<i>Baptisia tinctoria</i> , host plant of <i>Comandra</i>	
Inheritance of Fertility in Swine.....	1145-1160	<i>umbellata</i>	134
Automatic Transpiration Scale of Large		Barley. See <i>Hordeum</i> .	
Capacity for Use with Freely Exposed		Bartram, H. E. (paper), Effect of Natural	
Plants, An (paper).....	117-133	Low Temperature on Certain Fungi and	
Avena—		Bacteria.....	651-655
<i>chalar</i> , host plant of <i>Puccinia phleiprater-</i>		Bean—	
<i>sis</i>	212, 215	See <i>Phaseolus vulgaris</i> .	
<i>falsa</i> , host plant of <i>Puccinia phleipraterensis</i>	212-213	Florida velvet. See <i>Stizolobium doringi-</i>	
<i>sativa</i> —		<i>num</i> .	
absorption of boron by.....	884	Philippine Lyon. See <i>Stizolobium nigrum</i> .	
composition of seeds of.....	1162	Beef, mature—	
effect of—		comparison with immature veal.....	667-711
alkali salts on growth of.....	24ff	digestibility of.....	684-708
green manure on germination of.....	1165-1166	Beet, sugar. See <i>Beta vulgaris</i> .	
on germination of various seeds.....	1163-1166	Beetle—	
sulphur on the growth of.....	245-247	blister. See <i>Epicauta vittata</i> .	
host plant of <i>Puccinia phleipraterensis</i>	211-212	cherry leaf. See <i>Galerucella cavicollis</i> .	
method of correcting for yields of.....	1041,	Colorado potato. See <i>Leptinotarsa decem-</i>	
1045-1048		<i>lineata</i> .	
occurrence of manganese in.....	353	Belling, J. (paper), Inheritance of Length of	
resistance of, to alkali.....	22-24	Pod in Certain Crosses.....	405-420
transpiration of.....	592-597	Berg, W. N. (paper), Biochemical Compari-	
Avocado. See <i>Persea gratissima</i> .		sons between Mature Beef and Immature	
Bacillus—		Veal.....	667-711
<i>melonis</i> , effect of low temperature on.....	654	Beriberi and Cottonseed Poisoning in Pigs	
<i>tracheiphilus</i> , parasite of <i>Cucumis sativus</i> ...	259	(paper).....	489-493
<i>typhorus</i> , effect of low temperature on.....	654	<i>Beta vulgaris</i> —	
Back, E. A., and Pemberton, C. E. (paper)—		absorption of boron by.....	880
Banana as a Host Fruit of the Mediter-		effect of alkali salts on the growth of.....	4-6, 24ff
ranean Fruit Fly.....	793-804	factors influencing—	
Effect of Cold-Storage Temperatures upon		infection by <i>Cercospora beticola</i>	1029-1037
the Mediterranean Fruit Fly.....	657-666	stomatal movement in.....	1012-1019
Bacteria—		histological relations of <i>Phoma betae</i> to.....	55-58
effect of—		host plant of—	
natural low temperature on.....	651-655	<i>Cercospora beticola</i>	1011-1018
on germination.....	1171-1173	<i>Peronospora schachlii</i>	59-67
soil—		resistance of, to alkali.....	22-24
effect of—		Betula—	
calcium sulphate on.....	772-774	<i>nigra</i> , host plant of <i>Comandra umbellata</i>	134
sulphur on.....	772-774	<i>populifolia</i> , host plant of <i>Comandra um-</i>	
non-spore-forming.....	939	<i>bellata</i>	134
relation to crop-producing power.....	855-869	Biochemical Comparisons between Mature	
slime-forming.....	940	Beef and Immature Veal (paper).....	667-711
Bacterium—		Biology of <i>Apanteles militans</i> (paper).....	495-508
<i>abortus</i> : agglutination test for in milk.....	871-875	Bishop, E. S., and Alway, F. J. (paper), Ni-	
<i>lachrymans</i> —		trogen Content of the Humus of Arid Soils.....	909-916
causal organism of angular leafspot.....	466-474	Bituminous materials, test of, with new pen-	
control of.....	474-475	etration needle.....	1121-1126
geographical distribution of.....	467	Blisterbeetle. See <i>Epicauta vittata</i> .	
identification and isolation of.....	466-467	Boll weevil. See <i>Anthonomus grandis</i> .	
morphology and physiology of.....	470-474	Boron: Its Absorption and Distribution in	
		Plants and Its Effect on Growth (paper).....	87-890

	Page		Page
Bouyoucos, G. J. (paper), Effect of Temperature on Movement of Water Vapor and Capillary Moisture in Soils.....	141-172	<i>Castilleja miniata</i> , host plant of <i>Cronartium coleosporioides</i>	781-785
<i>Brassica</i> —		Castor bean. See <i>Ricinus communis</i> .	
<i>alba</i> —		Cations, effect on growth of <i>Triticum</i> spp.....	42-43
composition of seed of.....	1162	Cats, experiments in feeding beef and veal to.....	703-708
effect of green manure on germination of.....	1162	Cement, asphalt—	
<i>napus</i> , effect of sulphur on growth of.....	243-245	characteristics of.....	805-807
<i>nigra</i> , host plant of <i>Cystopus candidus</i>	63	effect of—	
Briggs, L. J., and Shantz, H. L. (paper)—		controllable variables upon the penetration test for.....	805-818
An Automatic Transpiration Scale of Large Capacity for Use with Freely Exposed Plants.....	217-232	load variations on.....	815-816
Hourly Transpiration Rate on Clear Days as Determined by Cyclic Environmental Factors.....	583-650	temperature variation on.....	813-815
<i>Bromus tectorum</i> , host plant <i>Puccinia phlebitidis</i>	212-233, 215	time variations on.....	816-817
Brown, P. E. (paper), Relation between Certain Bacterial Activities in Soils and Their Crop-Producing Power.....	855-869	methods of preparation for testing.....	807-813
Brownrot, varietal resistance of plums to.....	365-396	<i>Cephalothecium roseum</i> , effect of low temperature on.....	652
Bryan, M. K., and Smith, E. F., (paper), Angular Leafspot of Cucumbers.....	405-476	<i>Ceratitis capitata</i> —	
Buckner, G. Davis (paper), Translocation of Mineral Constituents of Seeds and Tubers of Certain Plants During Growth.....	449-458	infestation of <i>Musa</i> spp. by.....	793-804
Buckwheat. See <i>Fagopyrum tataricum</i> .		effect of cold-storage temperatures upon.....	657-666
Calcium—		parasite of—	
carbonate, effect of, on germination of seeds.....	1170	<i>Chrysophyllum cainito</i>	657
sulphate—		<i>Mangifera indica</i>	657
effect of—		<i>Musa</i> spp.....	793-804
on forms of plant life.....	771-780	<i>Persea gratissima</i>	657
on growth of higher plants.....	775-778	presence of, in banana plantation.....	793-794
on legume bacteria.....	774-775	<i>Cercospora</i> —	
Calorimeter, respiration—		<i>beticola</i> —	
air-purifying system in.....	305-310	causal organism of leafspot.....	1011
air-tension equalizer in.....	304-305	factors influencing infection by.....	1029-1037
description of.....	299-348	parasite of <i>Beta vulgaris</i>	1011-1038
improved.....	299-348	relation of stomatal movement to infection by.....	1011-1038
principle of.....	301-302	<i>personata</i> —	
Canal, irrigation—		casual organism of leafspot of <i>Arachis hypogaea</i>	891-902
computation of discharge of.....	226-231	control of.....	891-894
use of current meters in.....	217-232	damage caused by.....	894-895
<i>Cannabis sativa</i> —		dissemination of.....	895-901
composition of seed of.....	1163	parasite of <i>Arachis hypogaea</i>	891-902
effect of green manure on germination of.....	1162-1166	<i>Chauliognathus</i> sp., carrier of <i>Cercospora personata</i>	898-899
Capillary moisture in soils, effect of temperature on movement of.....	141-172	Cherry and Hawthorn Sawfly Leaf Miner (paper).....	519-528
<i>Capsella bursa pastoris</i> , host plant of <i>Cystopus candidus</i>	63, 67	Cherry leaf beetle. See <i>Galerucella caricollis</i> .	
Carbohydrate Transformations in Sweet Potatoes (paper).....	543-560	<i>Chimaphila umbellata</i> , host plant of <i>Comandra umbellata</i>	134
Carbon dioxide, effect of, on germination.....	1169	<i>Chrysophyllum cainito</i> , host fruit of <i>Ceratitis capitata</i>	657
<i>Carr</i> sp., host plant of <i>Comandra umbellata</i>	134	<i>Chrysopsis mariana</i> , host plant of <i>Comandra umbellata</i>	134
Carpenter, C. W. (paper), Some Potato Tuber-Rots Caused by Species of <i>Fusarium</i>	183-210	Climate, relation of, to insect development.....	1183-1185, 1190-1191
Carrero, J. O., and Gile, P. L. (paper), Ash Composition of Upland Rice at Various Stages of Growth.....	357-364	Clover—	
Carruth, F. E., and Withers, W. A. (paper), Gossypol, the Toxic Substance in Cottonseed Meal.....	261-283	red. See <i>Trifolium pratense</i> .	
<i>Castanea dentata</i> , host plant of <i>Comandra umbellata</i>	134	See <i>Trifolium</i> .	
		sweet. See <i>Melilotus alba</i> .	
		Cold-storage temperatures, effect of, upon the Mediterranean fruit fly.....	657-666
		Coleman, D. A., Lint, H. C., and Kopeloff, N. (paper), Separation of Soil Protozoa.....	137-140
		Coleoptera, agency in dissemination of <i>Cercospora personata</i>	898-901
		<i>Colletotrichum lindemuthianum</i> , effect of low temperature on.....	654

	Page		Page
<i>Comandra</i> —		<i>Cylindrosporium pomii</i> , effect of low tempera-	
<i>trida</i> , host plant of <i>Peridermium pyriforme</i>	133	ture on.....	652
<i>pallida</i> , host plant of <i>Peridermium pyriforme</i>	133	<i>Cyrtopus candidus</i> —	
<i>richardsonia</i> , host plant of <i>Peridermium pyriforme</i>	133	parasite of <i>Lepidium virginicum</i>	62-63, 67
<i>umbellata</i> —		perennial mycelium in.....	62-63, 67
host plant of <i>Peridermium pyriforme</i>	133-289	<i>Dactylis glomerata</i> —	
host plants of.....	134	host plant of <i>Puccinia phlepratensis</i>	211-215
parasitism of.....	133-135	<i>Danthonia compressa</i> , host plant of <i>Comandra umbellata</i>	134
Combination, new.....	1006	<i>Diabrotica vittata</i> , agent in spread of bacterial wilt.....	257
<i>Comptonia peregrina</i> , host plant of <i>Comandra umbellata</i>	134	Diascise—	
Concrete roads, apparatus for measuring the wear of.....	951-954	in apple flesh, results of tests for.....	100
Cone, V. M. (paper)—		presence of, in apple juice.....	158
A New Irrigation Weir.....	1127-1136	<i>Dipbacus fullonum</i> , host plant of <i>Peronospora dipsaci</i>	59-67
Flow through Weir Notches with Thin Edges and Full Contractions.....	1051-1113	Dissemination of Bacterial Wilt of Cucurbits (paper).....	257-260
Cook, F. C. (paper), Boron: Its Absorption and Distribution in Plants and Its Effect on Growth.....	877-890	Distribution of the Virus of the Mosaic Disease in Capsules, Filaments, Anthers, and Pistils of Affected Tobacco Plants (paper).....	251-256
Coolidge, L. H. (paper), Agglutination Test as a Means of Studying the Presence of <i>Bacterium abortus</i> in Milk.....	871-875	Diuresis and Milk Flow (paper).....	561-568
Coons, G. H. (paper), Factors Involved in the Growth and Pycnidium Formation of <i>Plenodomus fuscomaculans</i>	713-769	Diuretics, effect of, on milk flow of goats.....	561-567
Copper sulphate, effect on <i>Bacterium lactymani</i>	474-475	Dryrot—	
Corn. See <i>Zea mays</i> .		caused by—	
Cotton—		<i>Fusarium eumartii</i>	198-201
See <i>Gossypium</i> .		<i>Fusarium rodenticola</i>	195-196
wild. See <i>Thurberia thespisoides</i> .		Edson, H. A. (paper), Histological Relations of Sugar-Beet Seedlings and Phoma betae.....	55-58
Cottonseed—		Effect of Alkali Salts in Soils on the Germination and Growth of Crops (paper).....	1-53
kernels, toxicity of.....	265-266, 273-283	Effect of Cold-Storage Temperatures upon the Mediterranean Fruit Fly (paper).....	617-666
meal—		Effect of Controllable Variables upon the Penetration Test for Asphalts and Asphalt Cement (paper).....	805-818
comparison with rice as a feed for pigs.....	490-493	Effect of Elemental Sulphur and of Calcium Sulphate on Certain of the Higher and Lower Forms of Plant Life (paper).....	771-789
effect of, on breeding sows.....	491-492	Effect of Natural Low Temperature on Certain Fungi and Bacteria (paper).....	651-655
toxic substance in.....	261-288	Effect of Refrigeration upon the Larvae of <i>Trichinella spiralis</i> (paper).....	819-834
poisoning in pigs, relation of, to beriberi.....	489-493	Effect of Temperature on Movement of Water Vapor and Capillary Moisture in Soils (paper).....	141-172
toxicity of.....	261-262, 283-286	Egg production of domestic fowl, measurement of winter cycle in.....	439-437
Cowpea. See <i>Vigna sinensis</i> .		<i>Elymus</i> —	
<i>Crataegus</i> spp., host plants of <i>Profenusa col-laris</i>	520-521	<i>canadensis</i> , host plant of <i>Puccinia graminis aeneae</i>	215
<i>Cromartium</i> —		<i>robustus</i> , host plant of <i>Puccinia graminis aeneae</i>	215
<i>coleosporoides</i> , parasite of <i>Castilleja miniata</i>	781	<i>virginicus</i> , host plant of <i>Puccinia phlepratensis</i>	212-213, 215
<i>pyriforme</i> , uredinal form of <i>Peridermium pyriforme</i>	289	Emmer. See <i>Triticum dicoccum</i> .	
Crop, effect of alkali salts in soils on germination and growth of.....	1-53	Emulsin, determination of, in flesh of <i>Malus sylvestris</i>	111
Crosses, inheritance of length of pod in.....	405-420	Ectoparasite, caudal vesicle of.....	504-506
Cruciferae, effect of sulphur on.....	242-245	Environment—	
Cucumber. See <i>Cucumis sativus</i> .		hourly transpiration rate on clear days as determined by.....	583-590
<i>Cucumis sativus</i> —		relation of, to transpiration of plants.....	587-590
angular leafspot of.....	469-476		
host plant of—			
<i>Bacillus tracheiphilus</i>	250		
<i>Bacterium lactymani</i>	466-474		
<i>Plasmopara cubensis</i>	259		
Cucurbit, dissemination of bacterial wilt of.....	257-260		
<i>Cucurbita longpa</i> , absorption of boron by.....	879		
Current meter, use of, in irrigation canals.....	217-222		
Curtis, Maymie R. (paper), Frequency of Occurrence of Tumors in the Domestic Powl.....	397-404		
<i>Cypripedium puberulum</i> , host plant of <i>Rosellinia boreocensis</i>	1005		

	Page		Page
Enzymes of Apples and Their Relation to the Ripening Process (paper).....	103-116	<i>Fusarium</i> —Continued.....	
<i>Epicauta vittata</i> , carrier of <i>Cercospora personata</i>	898-899	<i>discolor</i> var. <i>sulphureum</i> —	
<i>Eriosoma pyri</i> , identity of.....	1115-1119	diagnostic characters of.....	207
Errata.....	VII	parasite of <i>Solanum tuberosum</i>	184-185
Esterase, determination of, in flesh of <i>Malus sylvestris</i>	111, 112	taxonomic arrangement of.....	207
Evaporation, ratio of transpiration to.....	634-637	<i>eumartii</i> —	
Experiments in the Use of Current Meters in Irrigation Canals (paper).....	217-222	cause of dryrot.....	198-201
Factors Influencing the Longevity of Soil Micro-organisms Subjected to Desiccation, with Special Reference to Soil Solution (paper).....	927-942	diagnostic characters of.....	203-205
Factors Involved in the Growth and Pycnidium Formation of <i>Penicillium fuscomaculans</i> (paper).....	713-769	taxonomic arrangement of.....	203-205
<i>Fagopyrum tataricum</i> —		<i>gibbosum</i> , parasite of <i>Solanum tuberosum</i>	184
composition of seeds of.....	1162	<i>hyperosysporum</i> —	
effect of green manure on germination of.....	1162	diagnostic characters of.....	206
Fate and Effect of Arsenic Applied as a Spray for Weeds (paper).....	459-463	parasite of <i>Ipomoea batatas</i>	189
Fertility, swine—		taxonomic arrangement of.....	206
inheritance of.....	1145-1160	<i>martii</i> , parasite of <i>Solanum tuberosum</i>	186, 201-203
nongenetic factors affecting.....	1146-1148	<i>moniliforme</i> , parasite of <i>Solanum tuberosum</i>	186, 201, 203
Fertilizer—		<i>orthoceras</i> , parasite of <i>Solanum tuberosum</i>	184
effect of, on North Carolina soils.....	577-581	<i>oxytrophum</i> , parasite of <i>Solanum tuberosum</i>	187-191
requirements, relation of, to petrography of North Carolina soil.....	567-582	<i>radicicola</i> —	
Field pea, Canada. See <i>Pisum arvense</i> .		cause of—	
Firefly. See <i>Chauliognathus</i> sp.		dryrot.....	195, 196
Flax. See <i>Linum usitatissimum</i> .		jelly-end rot.....	194-195
Float, surface, measurement of water velocity with.....	222-226	diagnostic characters of.....	205-206
Flow through Weir Notches with Thin Edges and Full Contractions (paper).....	1051-1113	taxonomic arrangement of.....	205-206
Forest nurseries, disease in, caused by <i>Peridermium filamentosum</i>	781-785	<i>rubiginosum</i> , syn. <i>Fusarium culmorum</i> .	
Fowl, domestic—		<i>solanii</i> —	
frequency of occurrence of tumors in.....	397-404	parasite of <i>Solanum tuberosum</i>	186, 188, 189, 201, 203
measurement of winter cycle in egg production of.....	429-437	diagnostic characters of.....	203-204
percentage of, affected with tumors.....	398	taxonomic arrangement of.....	203-204
<i>Fragaria</i> —		sp. of conifers, effect of low temperature on.....	652
<i>americana</i> , host plant of <i>Comandra umbellata</i>	134	spp. —	
<i>virginiana</i> , host plant of <i>Comandra umbellata</i>	134	results of inoculation of <i>Solanum tuberosum</i> with.....	201-203
Fred, E. B. (paper), Relation of Green Manures to Failure of Certain Seedlings.....	1161-1176	causal organism of potato tuber-rot.....	183-210
Frequency of Occurrence of Tumors in the Domestic Powl (paper).....	397-404	<i>subulatum</i> , parasite of <i>Solanum tuberosum</i>	184
Fungus—		<i>trichothecoides</i> —	
effect of—		diagnostic characters of.....	184, 185
natural low temperature on.....	651-655	parasite of <i>Solanum tuberosum</i>	207
on germination of seeds.....	1172-1174	taxonomic arrangement of.....	207
Further Studies on Peanut Leafspot (paper).....	891-902	<i>vasinfectum</i> , parasite of <i>Solanum tuberosum</i>	186, 192-194
<i>Fusarium</i> —		<i>ventricosum</i> , parasite of <i>Solanum tuberosum</i>	184
<i>coeruleum</i> —		<i>Galerucella caricis</i> —	
diagnostic characters of.....	203-204	control of.....	949
parasite of <i>Solanum tuberosum</i>	184	habits of.....	945-948
taxonomic arrangement of.....	203-204	life history of.....	945-948
<i>culmorum</i> , parasite of <i>Solanum tuberosum</i>	184	outbreaks.....	944
		parasite of <i>Prunus pennsylvanica</i>	944
		Gas, residual, determination of, in respiration calorimeter.....	310-314
		<i>Gaylussacia frondosa</i> , host plant of <i>Comandra umbellata</i>	134
		Germination—	
		effect of—	
		aeration on.....	1168
		ammonium hydroxid on.....	1168-1169
		bacteria on.....	1171-1172
		carbon dioxid on.....	1169
		fungi on.....	1172-1174
		green manure on.....	1161-1165
		heat on.....	1170-1171
		position of green manure on.....	1167
		soil type on.....	1167
		temperature on.....	1168
		time of planting on.....	1164-1165

	Page		Page
Cile, P. L., and Carrero, J. O. (paper), Ash Composition of Upland Rice at Various Stages of Growth.....	357-364	<i>Gymnosporangium</i> —Continued.	
Giltner, W., and Langworthy, H. V. (paper), Some Factors Influencing the Longevity of Soil Micro-organisms Subjected to Desiccation, with Special Reference to Soil Solution.....	927-942	<i>koreaense</i> —	
Ginseng. See <i>Panax quinquefolium</i> .		description of.....	1006-1007
<i>Glomerella rufomaculans</i> , effect of low temperature on.....	652	economic importance of.....	1007-1008
<i>Glycine</i> —		<i>phoiniae</i> —	
<i>hispida</i> , absorption of boron by.....	883-884	description of.....	1007
<i>soja</i> —		economic importance of.....	1008
composition of seeds of.....	1162	Harding, S. T. (paper), Experiments in the Use of Current Meters in Irrigation Canals.....	217-232
effect of green manure on germination of.....	1164-1166	Harris, Frank S. (paper), Effect of Alkali Salts in Soils on the Germination and Growth of Crops.....	1-53
Goat, effect of diuretics on milk flow of.....	561-567	Hart, E. B., and Tottingham, W. E. (paper), Relation of Sulphur Compounds to Plant Nutrition.....	233-250
Goldbeck, A. T. (paper), Apparatus for Measuring the Wear of Concrete Roads.....	951-954	Harter, L. L. (paper), Sweet-Potato Scurf.....	787-791
<i>Gossypium</i> —		Hasselbring, H., and Hawkins, L. A. (paper): Carbohydrate Transformations in Sweet Potatoes.....	543-556
<i>herbaceum</i> —		Respiration Experiments with Sweet Potatoes.....	509-517
composition of seeds of.....	1162	Hawkins, L. A., and Hasselbring, H. (paper): Carbohydrate Transformations in Sweet Potatoes.....	543-556
effect of—		Respiration Experiments with Sweet Potatoes.....	509-517
fungi on germination of.....	1172-1174	Hawthorn sawfly leaf miner. See <i>Proctosus collaris</i> .	
green manure on germination.....	1163-1170	Headden, W. P. (paper), Occurrence of Manganese in Wheat.....	349-355
heat on germination of.....	1170-1171	Heartrot—	
temperature on germination of.....	1168	honeycomb—	
time of planting on germination of.....	1164-1165	character of, in <i>Quercus alba</i>	412-424
spp.—		distribution of.....	427-428
toxicity of seeds of.....	261-262	in <i>Quercus</i> spp.....	421-425
yields of, in North Carolina soils.....	578-580	resemblance to other rots.....	424-425
<i>Gossypol</i> —		pocketed. See Honeycomb heartrot.	
acetate, crystalline, toxicity of.....	273-278	Hedgcock, G. G. (paper), Parasitism of Comandra umbellata.....	133-135
extract, toxicity of.....	266-267	Hedgcock, G. G., and Long, W. H. (paper), Two New Hosts for Peridermium pyri-forme.....	289-290
method of—		<i>Helianthus</i> —	
feeding.....	265, 267-270	<i>annuus</i> —	
preparation.....	262-263	composition of seeds of.....	1162
occurrence of.....	264-265	effect of green manure on germination of.....	1162
oxidized, toxicity of.....	278-281	<i>divaricatus</i> , host plant of <i>Plasmopara halstedii</i>	65, 66, 67
post-mortem observations of, effects of administering.....	270, 273-275	<i>Helophila unipunctata</i> , host of <i>Apanites militaris</i>	40
precipitated, toxicity of.....	271-272	<i>Heliothis obsoleta</i> , agency in dissemination of <i>Cercospora personata</i>	808-809
properties of.....	264-265	Hemiptera, agency in dissemination of <i>Cercospora personata</i>	808-809
<i>Gossypol</i> , the Toxic Substance in Cottonseed Meal (paper).....	261-288	Hemp. See <i>Cannabis sativa</i> .	
Gramineae, effect of sulphur on.....	245-247	<i>Hepatica acutiloba</i> , host plant of <i>Plasmopara pyramis</i>	67
Grape, wild. See also <i>Vitis cordifolia</i> .		Herrick, G. W., and Matheson, R. (paper), Observations on the Life History of the Cherry Leaf Beetle.....	943-950
Grasshopper, carrier of <i>Cercospora personata</i>	808-900	Heterogeneity, soil, in variety tests, method of correcting for.....	1039-1050
Green manure. See Manure, green.		Hibernation of <i>Phytophthora infestans</i> in the Irish Potato (paper).....	71-102
<i>Gymnosporangium</i> , an Asiatic species of, established in Oregon.....	1003-1010	<i>Hieracium venosum</i> , host plant of <i>Comandra umbellata</i>	134
<i>Gymnosporangium</i> —			
<i>arsaticum</i> , hypomyces, <i>Gymnosporangium</i>			
<i>koreaense</i> .			
<i>Blasdaleum</i> , economic importance of.....	1007-1008		
<i>chinense</i> , syn. <i>Gymnosporangium koreaense</i> .			
<i>Haraceum</i> —			
parasite of <i>Juniperus chinensis</i>	1005		
syn. <i>Gymnosporangium koreaense</i> .			
<i>japonicum</i> , syn. <i>Gymnosporangium koreaense</i> .			
<i>amie</i> .			
<i>juniperi-virginianae</i> , economic importance of.....	1007-1008		

	Page		Page
Histological Relations of Sugar-Beet Seedlings and Phoma betae (paper).....	55-58	<i>Ionactis linariifolius</i> , host plant of <i>Comandra umbellata</i>	134
Honeycomb heartrot—		<i>Ipomoea batatas</i> —	
character of, in <i>Quercus alba</i>	422-424	analyses of.....	545-557
control of.....	428	carbohydrate transformations in.....	543-560
distribution of.....	427-428	host plant of <i>Monilochaetes infusans</i>	787-791,
resemblance of, to other rots.....	424-425	995-1002	
Honeycomb Heartrot of Oaks Caused by <i>Stereum subpileatum</i> (paper).....	421-428	respiration experiments with.....	599-517
<i>Hordeum</i> —		scurl of.....	787-791
sp., occurrence of manganese in.....	353	storage experiments with.....	997
spp.—		variations in composition of.....	510-516
effect of alkali salts on growth of.....	2311	Irrigation—	
resistance of, to alkali.....	22-24	canal, experiments in use of current meters	
<i>vulpare</i> —		in.....	217-232
effect of sulphur on growth of.....	245	effect of, on arsenic in soil.....	460-461
test of inoculation with <i>Puccinia phleipra-lensis</i>	211-212	weir, a new.....	1127-1136
Hourly Transpiration Rate on Clear Days as Determined by Cyclic Environmental Factors (paper).....	583-650	Jackson, F. H., Jr. and Hubbard, P. (paper), Relation Between the Properties of Hardness and Toughness of Road-Building Rock.....	903-907
Houston, D. F., Announcement of Weekly Publication.....	i	Jackson, H. S. (paper), An Asiatic Species of Gymnosporangium Established in Oregon.....	1003-1010
Hubbard, P., and Jackson, F. H., jr. (paper), Relation Between the Properties of Hardness and Toughness of Road-Building Rock.....	903-907	Jelly-end rot, caused by <i>Fusarium radicola</i>	184-195
Hubbard, P., and Pritchard, F. P. (paper), Effect of Controllable Variables upon the Penetration Test for Asphalts and Asphalt Cements.....	805-818	Jensen, L., and Stakman, E. C. (paper), Infection Experiments with Timothy Rust.....	211-216
Hubert, E. R., and Weir, J. R. (paper), A Serious Disease in Forest Nurseries Caused by <i>Peridermium filamentosum</i>	781-785	Johnson, P. M., and Ballinger, A. M. (paper), Life-History Studies of the Colorado Potato Beetle.....	917-926
Humidity, relationship of, to insect development.....	1183-1191	<i>Juniperus chinensis</i> , host plant of <i>Gymnosporangium Haraeonium</i>	1005
Humus—		Kamani nut. See <i>Terminalia catappa</i> .	
of arid soils, nitrogen content of.....	909-916	Katydid, agency in dissemination of <i>Cercospora personata</i>	898, 900
relation of, to nitrogen in California soils.....	914-915	Koch, George P. (paper), Activity of Soil Protozoa.....	477-488
Identity of <i>Eriosoma pyri</i> (paper).....	1115-1119	Kopeloff, N., Lint, H., and Coleman, D. A. (paper), Separation of Soil Protozoa.....	137-140
Improved Respiration Calorimeter for Use in Experiments with Man, An (paper).....	299-348	<i>Lactuca sativa</i> , absorption of boron by.....	879-882
Inactivity, zone of, in insect development.....	1189-1190	Lady beetle. See <i>Megilla maculata</i> .	
Infection by <i>Cercospora belizicola</i> , relation of stomatal movement to.....	1011-1032	Langworthy, H. V., and Giltner, W. (paper), Some Factors Influencing the Longevity of Soil Micro-organisms Subjected to Desiccation, with Special Reference to Soil Solution.....	927-942
Infection Experiments with Timothy Rust (paper).....	211-216	<i>Laphygma frugiperda</i> , periods of activity of.....	1189
Influence of Growth of Cowpeas upon Some Physical, Chemical, and Biological Properties of Soil (paper).....	439-448	Leaf—	
Inheritance of Fertility in Swine (paper).....	1145-1160	hopper, carrier of <i>Cercospora personata</i>	898, 900
Inheritance of Length of Pod in Certain Crosses (paper).....	495-420	maturity, definition of.....	1011
Insect—		miner, sawfly. See <i>Profenusa collaris</i> .	
agency in dissemination of <i>Cercospora personata</i>	897-901	Leafspot—	
development—		angular, of <i>Cucumis sativus</i>	465-476
climatic relations in.....	1184-1185	injury to <i>Arachis hypogaea</i>	894-895
relationships of temperature and humidity to.....	1173-1181	occurrence on—	
nomenclature of climatic effects on life of.....	1190-1191	<i>Arachis hypogaea</i>	891-902
zone of inactivity in development of.....	1189-1190	<i>Beta vulgaris</i>	1011
Invertase in apple flesh, results of tests for.....	109-111	Le Clair, C. A. (paper), Influence of Growth of Cowpeas upon Some Physical, Chemical, and Biological Properties of Soil.....	439-448
		<i>Lepidium virginicum</i> —	
		host plant of—	
		<i>Cystolopus candidus</i>	67
		<i>Peronospora parastica</i>	60-63, 67

	Page		Page
Lepidoptera, agency in dissemination of		Massee, views on origin of <i>Phytophthora</i>	
<i>Cercospora personata</i>	898-901	<i>infestans</i>	96-97
<i>Leptinotarsa descendina</i> —		Matheson, R., and Herrick, G. W. (paper)	
eggs produced by.....	919-922	Observations on the Life History of the	
file history of.....	917-926	Cherry Leaf Beetle.....	941-950
<i>Lespedeza violacea</i> , host plant of <i>Comandra um-</i>		Meal, cottonseed, toxic substance in.....	261-288
bellata.....	134	Measurement of the Winter Cycle in the Egg	
Lettuce. See <i>Lactuca sativa</i> .		Production of Domestic Fowl (paper).....	429-437
Life-History Studies of the Colorado Potato		<i>Medicago sativa</i> —	
Beetle (paper).....	917-926	effect of alkali salts on growth of.....	25ff
Lint, H. C., Kopeloff, N., and Coleman, D. A.		resistance of, to alkali.....	22-24
(paper), Separation of Soil Protozoa.....	137-140	use of, as a diuretic.....	361
<i>Linum usitatissimum</i> , composition of seed of.....	1162	Mediterranean fruit fly. See <i>Ceratitis capi-</i>	
<i>Liquidambar styraciflua</i> , host plant of <i>Stereum</i>		tata.	
subpileatum.....	427	<i>Megilla maculata</i> , agency in dissemination of	
Litter, inheritance of size of, studies in.....	1149-1150	<i>Cercospora personata</i>	898-900
Lodgepole pine. See <i>Pinus murroyma</i> .		<i>Meibomia paniculata</i> , host plant of <i>Comandra</i>	
<i>Lolium</i> —		<i>umbellata</i>	134
<i>italicum</i> , host plant of <i>Puccinia phleipraten-</i>		Melhus, I. E. (paper)—	
sis.....	212-213, 215	Hibernation of <i>Phytophthora infestans</i> in	
<i>perenne</i> , host plant of <i>Puccinia phleipraten-</i>		the Irish Potato.....	71-102
sis.....	212-213, 215	Perennial Mycelium in Species of Perono-	
Long, W. H. (paper), Honeycomb Heartrot		sporaceae Related to <i>Phytophthora in-</i>	
of Oaks Caused by <i>Stereum subpileatum</i>	421-428	festans.....	59-70
Long, W. H., and Hedgcock, G. G. (paper),		Meter, current, experiments in use in irriga-	
Two New Hosts for <i>Peridermium pyri-</i>		tion canals.....	217-231
forme.....	289-290	Method of Correcting for Soil Heterogeneity	
Lupine, white. See <i>Lupinus albus</i> .		in Variety Tests, A (paper).....	1039-1050
<i>Lupinus albus</i> —		Micro-organism, soil, factors influencing the	
composition of seeds of.....	1162	longevity of.....	927-942
effects of green manure on germination of.....	1162	Milk—	
<i>Lycopersicon esculentum</i> , absorption of boron		agglutination test for presence of <i>Bacterium</i>	
by.....	881-882	<i>abortus</i> in.....	871-875
Lyon bean. See <i>Shizobium nigrum</i> .		flow, effect of diuresis on.....	561-568
<i>Lysimachia quadrifolia</i> , host plant of <i>Coman-</i>		Milner, R. D., and Langworthy, C. F. (paper),	
<i>umbellata</i>	134	An Improved Respiration Calorimeter for	
McGeorge, W. T. (paper), Fate and Effect of		Use in Experiments with Man.....	299-348
Arsenic Applied as a Spray for Weeds.....	459-463	Miner, leaf, sawfly. See <i>Protenusa collaris</i> .	
McKay, M. B., and Pool, V. W. (paper), Re-		Mineral constituents, translocation of, in	
lation of Stomatal Movement to Infection		seeds and tubers of plants.....	449-458
by <i>Cercospora beticola</i>	1011-1018	Moisture—	
Magnesium salts, effect of, on plant growth.....	5-6, 8-10, 16ff	and temperature, relation of, to spread of	
<i>Malus</i> spp.—		mycelium of <i>Phytophthora infestans</i> in	
changes in chemical composition during		tubers.....	73
ripening.....	105-106	capillary, effect of temperature on move-	
host fruit of <i>Ceratitis capitata</i>	659	ment of, in soils.....	141-172
host plant of—		movement of, between warm and cold	
<i>Monilia fructigena</i>	365-367	soils.....	141-172
<i>Plenodomus fuscomaculans</i>	713	soil, description of apparatus for determin-	
<i>Sclerotinia</i> spp.....	365	ing translocation of.....	147-144
relation of enzymes to ripening process of.....	105-110	<i>Monilia</i> —	
Manganese, occurrence of, in wheat.....	349-355	<i>fructigena</i> , parasite of <i>Malus</i> spp.....	361, 367
<i>Mangifera indica</i> , host fruit of <i>Ceratitis capi-</i>		<i>laxa</i> , parasite of apricot.....	365
tata.....	657	<i>Monilochaetes</i> —	
Mango. See <i>Mangifera indica</i> .		<i>infusans</i> —	
Manure, green—		causal organism of soil stain.....	998
effect of, on seed germination.....	1162-1165	description of.....	789-791, 1000
nature of injurious agent in.....	1166	distribution of.....	797
relation of—		inoculation experiments with.....	788-790
to failure of seedlings.....	1162-1176	isolation of.....	788
history of oil seeds.....	1174	loss due to.....	788
<i>Maple</i> —		morphology of.....	998-1000
sugar. See <i>Acer saccharum</i> .		parasite of <i>Ipomoea batatas</i>	787-791, 993-1002
water. See <i>Acer saccharum</i> .		physiology of.....	998-1000
		taxonomy of.....	790-791, 10000
		technical description of genus.....	797

- Morphology and Biology of the Green Apple
Aphis (paper)..... 955-994
- Mosaic disease—
distribution of the virus of..... 251-256
occurrence of virus in plants of *Nicotiana
glabrum*..... 254
virus of, in placental structure of plants of
Nicotiana glabrum..... 252-253
- Movement of moisture from warm to cold
soil..... 141-156
- Mucor* sp., parasite of *Solanum tuberosum*... 186,
201-203
- Musa* spp.—
green fruits of, immunity of, to attack by
Ceratitis capitata..... 799-801
immunity of, to attack by *Ceratitis capitata* 793-804
- Muscular work, apparatus for measuring... 347-343
- Mustard, white. See *Brassica alba*.
- Needle, penetration—
use of, in testing bituminous materials. 1121-1126
results of standardization test of..... 1123
results of tests of..... 1124-1126
- New Interpretation of the Relationships of
Temperature and Humidity to Insect De-
velopment, A (paper)..... 1183-1191
- New Irrigation Weir, A (paper)..... 1127-1136
- New Penetration Needle for Use in Testing
Bituminous Materials, A (paper)..... 1121-1126
- New Spray Nozzle, A (paper)..... 1177-1182
- Nicotiana glabrum*, distribution of the virus
of the mosaic disease in..... 252-256
- Nitrate production—
effect of—
calcium sulphate on..... 774-775
sulphur on..... 774-775
- Nitrogen Content of the Humus of Arid Soils
(paper)..... 900-916
- Nitrogen—
humus, methods for determining..... 911-913
relation of, to humus in California soils... 914-915
- North Carolina, petrography of soils of.... 567-577
- Nozzle, spray, a new..... 1177-1182
- Nursery, forest, disease in, caused by *Peri-
dermium filamentosum*..... 781-785
- Nutrition, plant, relation of sulphur com-
pounds to..... 233-250
- Oak, white. See *Quercus alba*.
- Oats. See *Avena sativa*.
- Observations on Life History of the Cherry
Leaf Beetle (paper)..... 943-950
- Occurrence of Manganese in Wheat (paper). 349-355
- Oospora scabies*, syn. *Actinomyces chromogenus*.
- Orthoptera, agency in dissemination of *Cer-
atosporea personata*..... 898-901
- Oryza sativa*, ash analyses of..... 359-363
- Oskamp, J. (paper), Soil Temperatures as
Influenced by Cultural Methods..... 173-179
- Oxidases, determination of, in flesh of *Malus
sylvestris*..... 112-113
- Panax quinquefolium*—
blackrot of..... 294-296
host plant of—
Alternaria panax..... 181-182
Phytophthora cactorum..... 59, 67
Sclerotinia libertiana..... 291
- Panax quinquefolium*—Continued. Page
inoculation of, with *Sclerotinia libertiana*
from various sources..... 292-293
pathogenicity and identity of *Sclerotinia
libertiana* and *Sclerotinia smilacina* on..... 291-298
rootrot of..... 181-182
whiterot of..... 291-294
- Panicum* sp., host plant of *Comandra umbel-
lata*..... 134
- Parasitism of *Comandra umbellata* (paper). 133-135
- Pathogenicity and Identity of *Sclerotinia
libertiana* and *Sclerotinia smilacina* on *Gin-
seng* (paper)..... 291-298
- Pea—
See *Pisum sativum*.
Canada field. See *Pisum arvense*.
- Peanut. See *Arachis hypogaea*.
- Pearl, R. (paper), Measurement of the Winter
Cycle in the Egg Production of Domestic
Fowl..... 429-437
- Pearl, R., and Surface, F. M. (paper), A
Method of Correcting for Soil Heterogeneity
in Variety Tests..... 1039-1050
- Pectinase, determination of, in flesh of *Malus
sylvestris*..... 112-113
- Pemberton, C. E., and Back, E. A. (paper)—
Banana as a Host Fruit of the Mediterrane-
an Fruit Fly..... 793-804
Effect of Cold-Storage Temperatures upon
Mediterranean Fruit Fly..... 657-666
- Penetration test, asphalt, effect of controllable
variables upon..... 805-818
- Perennial Mycelium in Species of *Peronospor-
aceae* Related to *Phytophthora infestans*
(paper)..... 59-70
- Peridermium*—
ceruleum, parasite of *Pinus dicaricata*..... 289
complanata, parasite of *Pinus (murrayana)
contorta*..... 290
filamentosum—
causal organism of disease in forest nur-
series..... 781-785
parasite of—
Castilleja miniata..... 781-785
Pinus contorta..... 783
Pinus ponderosa..... 781
montanum, parasite of *Pinus murrayana*... 785
- pyriforme*—
aerial form of *Cronartium pyriforme*..... 289
parasite of—
Comandra umbellata..... 289
Pinus arizonica..... 290
Pinus dicaricata..... 133, 289-290
Pinus (murrayana) contorta..... 133, 289-290
Pinus ponderosa..... 133
Pinus ponderosa scopulorum..... 133, 290
Pinus pungens..... 133, 290
Pinus rigida..... 133, 289-290
two new hosts for..... 289-290
spp., parasites of *Quercus* spp..... 783
- Peronosporaceae*, relation of perennial myce-
lium in, to *Phytophthora infestans*..... 59-70
- Peronospora*—
alsinacrum, parasite of *Stellaria media*... 59, 67, 68
dipsaci, parasite of *Dipsacus fullonum*..... 59, 67
effusa—
parasite of—
Atriplex hortensis..... 59, 67
Spinacia oleracea..... 59, 67, 68

	Page		Page
Pritchard, F. P., and Reeve, C. S. (paper), A New Penetration Needle for Use in Testing Bituminous Materials.....	1121-1126	<i>Quercus</i> —Continued.	
<i>Prociophilus</i> —		<i>digitata</i> , host plant of <i>Comandra umbellata</i> ...	134
<i>aceris</i> , description of.....	1118	<i>marilandica</i> , host plant of <i>Comandra umbellata</i>	134
<i>alnifoliae</i> , description of.....	1118	<i>nana</i> , host plant of <i>Comandra umbellata</i>	134
<i>humulæ</i> , description of.....	1118	spp.—	
<i>corrugulans</i> , description of.....	1118	honeycomb heartrot of.....	421-428
<i>fragmi-depetalæ</i> , syn: <i>Prociophilus venafuscus</i>		host plant of—	
<i>fuscus</i>		<i>Peridermium</i> spp.....	783
<i>imbricator</i> , description of.....	1118	<i>Polyporus</i> spp.....	421
<i>populiconduplifolius</i> , description of.....	1118	<i>Stereum subpileatum</i>	421-427-428
<i>poschingeri</i> , description of.....	1119	Radish. See <i>Raphanus sativus</i> .	
<i>pyri</i> , description of.....	1116, 1118	Rand, F. V. (paper), Dissemination of Bacterial Wilt of Cucurbits.....	257-260
<i>tessellatus</i> , description of.....	1119	Ransom, B. H. (paper), Effects of Refrigeration upon the Larvæ of <i>Trichinella spiralis</i>	819-854
<i>venafuscus</i> , description of.....	1119	<i>Ranunculus</i> —	
<i>zyloiei</i> , description of.....	1119	<i>fascicularis</i> , host plant of <i>Peronospora ficariae</i>	59-64, 67
<i>Profenusa collaris</i> —		<i>ficaria</i> , host plant of <i>Peronospora ficariae</i>	64, 67
control of.....	527-528	Rape. See <i>Brassica napus</i> .	
description of.....	522-524	<i>Raphanus sativus</i> —	
distribution of.....	520-522	absorption of boron by.....	885
enemies of.....	526-527	effect of sulphur on growth of.....	242-243
habits of.....	524-526	Record, autographic transpiration.....	128-130
injury to <i>Prunus</i> spp. by.....	521-522	Reeve, C. S., and Pritchard, F. P. (paper), A New Penetration Needle for Use in Testing Bituminous Materials.....	1121-1126
life history of.....	524-526	Refrigeration, effect of, upon larvæ of <i>Trichinella spiralis</i>	819-854
parasite of—		Relation between Certain Bacterial Activities in Soils and Their Crop-Producing Power (paper).....	855-869
<i>Crataegus</i> spp.....	519-520	Relation between the Properties of Hardness and Toughness of Road-Building Rock (paper).....	903-907
<i>Prunus</i> spp.....	519-520	Relation of Green Manures to the Failure of Certain Seedlings (paper).....	1161-1176
Protease, determination of, in flesh of <i>Malus sylvestris</i>	113-114	Relation of Stomatal Movement to Infection by <i>Cercospora beticola</i> (paper).....	1012-1018
Protozoa, soil—		Relation of Sulphur Compounds to Plant Nutrition (paper).....	213-250
activity of.....	477-488	Resistance, varietal, of plums to brownrot..	365-396
encystment of.....	485-487	Respiration—	
separation of.....	137-140	calorimeter—	
<i>Prunus</i> —		conditions affecting and measurement of heat in.....	315-347
<i>pennsylvanica</i> , host plant of <i>Galerucella corticollis</i>	944	improved.....	399-348
spp.—		test of accuracy of.....	344-346
host plant of—		chamber, construction of.....	302-304
<i>Monilia</i> spp.....	365	Respiration Experiments with Sweet Potatoes (paper).....	509-517
<i>Profenusa collaris</i>	519-520	Respiratory exchange in respiration chamber, determination of.....	304-314
<i>Sclerotinia</i> spp.....	365-366	<i>Rhizoctonia</i> sp.—	
hybrids used in experiments with brownrot.....	369-370	effect of, on germination.....	1173-1174
relation of tannin content of, to resistance to <i>Sclerotinia</i> spp.....	389-390	parasite of <i>Solanum tuberosum</i>	186, 201, 202
resistance of, to brownrot.....	369, 379-383, 387	<i>Rhizopus nigricans</i> , parasite of <i>Solanum tuberosum</i>	183
susceptibility of, to brownrot.....	387	<i>Rhus copallina</i> , host plant of <i>Comandra umbellata</i>	134
varietal resistance of, to brownrot.....	365-396	Rice—	
<i>Pseudomonas</i> —		polished, effect of feeding to pigs.....	490-492
<i>campestris</i> , effect of low temperature on.....	654	See also <i>Oryza sativa</i> .	
<i>radicola</i> , longevity of, under varying conditions.....	932-940	upland, ash composition of, at various stages of growth.....	357
<i>Puccinia</i> —		<i>Ricinus communis</i> , composition of seed of... ..	1162
<i>graminis</i> —			
<i>avenae</i> , parasite of cereals and grasses... ..	213, 215		
<i>hordei</i> , parasite of grasses and cereals... ..	213, 215		
<i>phlebotensis</i> , infection experiments with.....	211-216		
Pycnidium formation in <i>Pseudomonas fuscomaculans</i>	713-769		
<i>Pyrus</i> —			
<i>communis</i> , host plant of <i>Roestelia koreensis</i>	1005		
<i>sinensis</i> , host plant of <i>Roestelia koreensis</i>	1005-1006		
<i>Quercus</i> —			
<i>alba</i> , character of honeycomb heartrot in.....	422-424		
<i>coccinea</i> , host plant of <i>Comandra umbellata</i>	134		

	Page		Page
Ripero, occurrence of, in <i>Prunus</i> spp.	388-389	<i>Sclerotinia</i> —	
Road, concrete, apparatus for measuring the wear of	951-954	<i>cinerea</i> —	
Roasting-ear worm. See <i>Heliothis obsoleta</i> .		effect of low temperature on	652
Rock, road-building—		parasite of <i>Prunus</i> spp.	365-366
relation between the properties of hardness and toughness of	903-907	toxicity of fruit acids to	388
tests of	903-907	<i>fructigena</i> , parasite of <i>Prunus</i> spp.	366
<i>Roestelia</i> —		<i>lisa</i> , parasite of <i>Prunus</i> spp.	365-366
<i>koreaensis</i> —		<i>libertiana</i> —	
occurrence of—		identity of	291-298
in America	1004-1006	pathogenicity of	291-298
in Japan	1004	<i>smilacina</i> —	
in Oregon	1005-1006	identity of	291-298
parasite of—		pathogenicity of	291-298
<i>Cydonia vulgaris</i>	1005	spp.—	
<i>Pyrus sinensis</i>	1003-1006	parasite of <i>Malus</i> spp.	365
relation of, to <i>Gymnosporangium koreanse</i>	1004-1006	pathological relations of	374-396
<i>pholimae</i> , syn. <i>Gymnosporangium pholimae</i> .		physiological relations	374-383
Rommel, G. M., and Vedder, E. B. (paper).	459	relation of tannin content of, to resistance of <i>Prunus</i> spp.	379-396
Beriberi and Cottonseed Poisoning in Pigs.	493	taxonomy of	379-384
Root—		Scurf of <i>Ipomoea batatas</i> —	
<i>let.</i> , occurrence of, in <i>Panax quinquefolium</i>	181-182	appearance of	787
<i>Rosa</i> —		injury caused by	787-788
<i>blanda</i> , host plant of <i>Comandra umbellata</i>	134	occurrence of	965-1002
<i>canina</i> , host plant of <i>Comandra umbellata</i>	134	distribution of	787
Rosenbaum, J. (paper). Pathogenicity and Identity of <i>Sclerotinia libertiana</i> and <i>Sclerotinia smilacina</i> on Ginseng	291-298	<i>Secale cereale</i> —	
Rosenbaum, J., and Zinnmeister, C. L. (paper). Alternaria panax, the Cause of a Root-rot of Ginseng	181-182	host plant of <i>Puccinia phleipratensis</i>	211-212
Rot—		occurrence of manganese in	353
brown, varietal resistance of plums to	365-366	transpiration of	602-609
dry, caused by <i>Fusarium</i> spp.	195-201	Seed—	
field and storage, occurrence of, in <i>Solanum tuberosum</i>	187-201	effect of green manures on germination of	1161-1165
heart—		oil, relation of green manure to injury of	1172
honeycomb—		Separation of Soil Protozoa (paper)	137-140
distribution of	427-428	Serious Disease in Forest Nurseries Caused by <i>Peridermium filamentosum</i> , A (paper)	781-785
occurrence of, in <i>Quercus</i> spp.	421-425	Shantz, H. L., and Briggs, L. J. (paper)—	
resemblance to other rots	424-425	An Automatic Transpiration Scale of Large Capacity for Use with Freely Exposed Plants	117-132
pocketed. See Honeycomb heartrot.		Hourly Transpiration Rate on Clear Days as Determined by Cyclic Environmental Factors	583-630
jelly-end, caused by <i>Fusarium radicola</i>	104-105	Shedd, O. M. (paper). Variations in Mineral Composition of Sap, Leaves, and Stems of the Wild-Grape Vine and Sugar-Maple Tree	529-542
ripe, occurrence of, in <i>Prunus</i> spp.	388-389	<i>Sisymbrium officinale</i> , host plant of <i>Cytophus candidus</i>	63
root, occurrence of, in <i>Panax quinquefolium</i>	181-182	<i>Smilacina racemosa</i> , results of inoculation with <i>Sclerotinia smilacina</i>	295-296
white, occurrence of, in <i>Panax quinquefolium</i>	201-204	Smith, E. F., and Bryan, M. K. (paper). Angular Leafspot of Cucumbers	468-478
<i>Rubus</i> —		Sodium—	
<i>canadensis</i> , host plant of <i>Comandra umbellata</i>	134	effect of, on milk flow	563-565
<i>procumbens</i> , host plant of <i>Comandra umbellata</i>	134	arsenite, effect of, on plant growth	459-460
<i>villosus</i> , host plant of <i>Comandra umbellata</i>	134	salts, effect of, on plant growth	510, 1551
<i>Rumex acetosa</i> , host plant of <i>Pernatospora rumicis</i>	67	Soil—	
Rust, timothy. See <i>Puccinia phleipratensis</i> .		arid, nitrogen content of humus of	999-1016
Rye. See <i>Secale cereale</i> .		bacteria—	
		effect of calcium sulphate on	772-774
		effect of sulphur on	772-774
		factors influencing longevity of	927-942
		non-spore-forming	939
		relation of, to crop-producing power	855-866
		slime-forming	940
		moisture, description of apparatus for determining translocation of	142-144
Salt, alkali, in soils, effect of, on germination and growth of crops	1-53		
<i>Scutellaria palmarum</i> , <i>Stx. Profenusa collaris</i> .			
transpiration, description	117-132		

- Soil—Continued.** Page
 conditions, effect of, on protozoa..... 479-485
 effect of—
 alkali salts in, on germination and growth
 of crops..... 1-53
Vigna sinensis on properties of..... 443-447
 heterogeneity of, in variety tests, method
 of correcting for..... 1039-1050
 North Carolina—
 mineralogical composition of..... 571-580
 relation of petrography of, to fertilizer re-
 quirements..... 569-582
 yields of *Gossypium* spp. on..... 578-580
 Utah, analysis of..... 14
 protozoa—
 activity of..... 477-488
 separation of..... 137-140
 properties of, influence of growth of *Vigna*
sinensis on..... 439-448
 type of, effect on germination..... 1167
Soil Temperatures as Influenced by Cultural
Methods (paper)..... 173-179
Soilstain—
 causal organism of..... 998
 effect of, on *Ipomoea batatas*..... 990
 factors favorable to development of..... 990
 symptoms of..... 996
Soilstain, or Scurf, of the Sweet Potato
(paper)..... 995-1002
Soja var. See *Glycine*.
Solanum—
jasminoides, host plant of *Leptimorpha de-*
clineolata..... 917
tuberosum—
 absorption of boron by..... 879-884
 analysis of..... 457-458
 growth of mycelium of *Phytophthora in-*
festans in..... 74-80
 hibernation of *Phytophthora infestans* in..... 71-102
 host plant of—
Fusarium spp..... 183-203
Leptimorpha declivata..... 917-926
Phytophthora infestans..... 59, 66, 67, 71-102, 183
Rhizopus nigricans..... 183
Sporotrichum flavissimum..... 186
Verticillium albo-atrum..... 186
 infection of, by conidia of *Phytophthora*
infestans..... 85-87
 inoculation of, with *Fusarium* spp..... 201-203
 relation of progeny of, to *Phytophthora in-*
festans in parent tuber..... 93-96
 translocation of mineral constituents of..... 457-458
Solar radiation, relation of, to transpiration. 631-634
Solidago—
bicolor, host plant of *Comandra umbellata*... 134
caesia, host of plant of *Comandra umbellata*... 134
juncea, host plant of *Comandra umbellata*... 134
memoralis, host plant of *Comandra umbellata*... 134
speciosa, host plant of *Comandra umbellata*... 134
Some Potato Tuber-Rots Caused by Species
of *Fusarium* (paper)..... 183-220
Sorghum. See *Andropogon sorghum*.
Soybean. See *Glycine*.
Species, new..... 204, 466
Sphaeropsis malorum, effect of low tempera-
 ture on..... 652
Spinacia oleracea, host plant of *Peronospora*
effusa..... 59, 67
Spiraea salicifolia, host plant of *Comandra*
umbellata..... 134
 Spore, resting, of *Phytophthora infestans*..... 98-99
Sporotrichum flavissimum, parasite of *Sola-*
num tuberosum..... 186, 201-202
Spray—
 advantage of fineness in..... 1179-1180
 advantage of uniformity in..... 1179-1180
 flat, advantages of..... 1179
 production of..... 1177-1179
 variations in fineness of..... 1180
 nozzle, new..... 1177-1182
 adjustment of..... 1181
 use of..... 1181
 Stakman, E. C., and Jensen, L. (paper) *Infection*
Experiments with Timothy Rust..... 211-216
 Star-apple. See *Chrysophyllum cainito*.
 Steenbock, H. (paper), *Diuresis and Milk*
Flow..... 561-563
Stellaria media, host plant of *Peronospora*
alsinearum..... 59, 67, 68
Stereum subpileatum—
 causal organism of honeycomb heartrot..... 421-423
 description of sporophore of..... 425
 distribution of..... 427-428
Sizolobium—
deeringianum, effect of crossing on length of
 pod..... 495-496
niveum, effect of crossing on length of pod..... 495-496
Stoma, movement of—
 factors influencing..... 1012-1022
 relation of, to infection by *Cercospora beti-*
cola..... 1011-1013
Sugar—
 beet. See *Beta vulgaris*.
 effect of—
 on ammonification..... 1169
 on carbon-dioxide evolution..... 1169-1172
 maple. See *Acer saccharum*.
Sulphur—
 effect of, on growth of plants..... 771-781
 compounds, relation of, to plant nutri-
 tion..... 233-250
Sunflower. See *Helianthus annuus*.
 Surface, F. M., and Pearl, R. (paper), *A*
Method of Correcting for Soil Heterogeneity
in Variety Tests..... 1039-1052
 Surface floats, measurement of water velocity
 with..... 222-225
 Sweet potato. See *Ipomoea batatas*.
 Sweet-Potato Scurf (paper)..... 787-791
Swine—
 deviations per generation in size of litter
 in..... 1150, 1151
 fertility in, nongenetic factors affecting..... 1146-1148
 value of herdbook data..... 1148-1151
 individual evidences of segregation in..... 1151-1155
 inheritance of fertility in..... 1145-1150
 litter frequency in..... 1149-1153
 Tannase, determination of, in *Malus sylves-*
tris..... 111
 Taubenhaus, J. J. (paper), *Soilstain, or Scurf,*
of the Sweet Potato..... 995-1002
Temperature—
 cold-storage, effect of, on *Ceratitis capitata*..... 657-666
 effect of, on movement of water and vapor,
 and capillary moisture in soils..... 141-147
 effective, in insect development..... 1185-1196

Temperature—Continued.	Page	Triticum—Continued.	Page
effect of, on germination of plants.....	2168	spp.—Continued.	
influence of, on growth of mycelium of <i>Phytophthora infestans</i>	77-80	method of correcting for yield of.....	1042,
low effect of, on certain fungi and bacteria.....	651-655	occurrence of—	1044-1045
relation of—		iron in.....	353
to insect development.....	1183-1191	manganese in.....	349-355
to spread of mycelium of <i>Phytophthora infestans</i> in <i>Solanum tuberosum</i>	73	resistance of, to alkali.....	22-24
soil—		vulgaris, inoculation of, with <i>Puccinia phlei-pratenensis</i>	217-212
as influenced by cultural methods.....	173-179	Tuber-rot, occurrence of, <i>Solanum tuberosum</i>	183-210
effect of different cultural methods.....	174-179	Tumors in domestic fowl—	
<i>Terminalia catappa</i> , host fruit of <i>Ceratitis capitata</i>	659	frequency of occurrence of.....	397-404
Thatcher, R. W. (paper). Enzymes of Apples and Their Relation to the Ripening Process.....	103-116	structure and location of.....	400-403
<i>Thurberia thespioidea</i> , host plant of <i>Anthonomus grandis</i>	1189-1190	Turmeric. See <i>Curcuma longa</i> .	
Timothy rust. See <i>Puccinia phlei-pratenensis</i> .		Turner, W. F., and Baker, A. C. (paper), Morphology and Biology of the Green Apple Aphis.....	955-994
Tomato. See <i>Lycopersicon esculentum</i> .		Two New Hosts for <i>Peridermium pyramiforme</i> (paper).....	289-290
Tottingham, W. E., and Hart, E. B. (paper). Relation of Sulphur Compounds to Plant Nutrition.....	233-250	Urea, effect of, on milk flow.....	563-566
Tow, Daniel G. (paper), Biology of <i>Apanetes militaris</i>	495-508	<i>Vaccinium—</i>	
Translocation of Mineral Constituents of Seeds and Tubers of Certain Plants During Growth (paper).....	449-458	<i>atrocarpum</i> , host plant of <i>Comandra umbellata</i>	134
Transpiration—		<i>nigrum</i> , host plant of <i>Comandra umbellata</i>	134
plant, measurement of.....	583-585	<i>racillans</i> , host plant of <i>Comandra umbellata</i>	134
relation of—		Valleau, W. D. (paper), Varietal Resistance of Plums to Brown-Rot.....	365-396
to environmental factors.....	637-646	Vapor, water, effect of temperature on movement of, in soils.....	141-172
to evaporation.....	634-637	Variations in Mineral Composition of Sap, Leaves, and Stems of the Wild-Grape Vine and Sugar-Maple Tree (paper).....	529-542
to solar radiation.....	631-634	Varietal Resistance of Plums to Brown-Rot (paper).....	365-396
graphs, comparison of.....	627-631	Variety tests, a method of correcting for soil heterogeneity in.....	1033-1050
rate, hourly, on clear days as determined by cyclic environmental factors.....	583-650	Veal, immature—	
scale, automatic, for use with freely exposed plants.....	117-132	comparison with mature beef.....	667-711
<i>Tremella koreanae</i> . Syn. <i>Gymnosporangium koreanae</i> .		digestibility of.....	684-708
Trichinae. See <i>Trichinella spiralis</i> .		Vedder, E. B., and Rommel, G. M. (paper), Beriberi and Cottonseed Poisoning in Pigs.....	489-493
<i>Trichinella spiralis—</i>		Velvet bean, Florida. See <i>Stizobium deerlingianum</i> .	
effect of—		<i>Venturia inaequalis</i> , effect of low temperature on.....	652
artificial digestion upon.....	847-850	<i>Veronica lederaefolia</i> , host plant of <i>Peronospora grisea</i>	59, 67, 68
refrigeration upon larvae of.....	819-854	Verticals, effect of varying numbers of, on the accuracy of current meter gaugings.....	226-231
refrigeration upon vitality of.....	817-845	<i>Verticillium albo-atrum</i> , parasite of <i>Solanum tuberosum</i>	386, 201-203
variations in vitality of.....	845	<i>Vicia sepium</i> , host plant of <i>Peronospora tritici</i>	59, 64, 65, 67
<i>Trichogramma minutum</i> , parasite of <i>Prodenia collaris</i>	516-527	<i>Vigna sinensis—</i>	
<i>Trifolium—</i>		absorption of boron by.....	881
<i>pratense</i> , effect of sulphur on growth of....	239-	effect of growth of, upon soil.....	439-448
	241, 778-779	influence of growth of, upon properties of soil.....	439-448
spp.—		<i>Vitis—</i>	
effect of—		<i>cordifolia—</i>	
on ammonification.....	1169	composition of—	
on carbon-dioxid evolution.....	1169-1170	leaves.....	536-538
on germination of various seeds.....	1161-1166	sap of.....	539-535
<i>Triticum—</i>		stems of.....	539-538
<i>aestivum</i> , effect of alkali salts on growth of..	4-11	variations in mineral composition of.....	529-542
<i>dicoccum</i> , occurrence of manganese in.....	353	<i>vinifera</i> , host plant of <i>Plasmopora viticola</i>	67
<i>sativum—</i>			
composition of seeds of.....	1163		
effect of green manure on germination of....	1163		
spp.—			
absorption of boron by.....	880, 883		

	Page		Page
maple. See <i>Acer saccharinum</i> .		Weir—Continued.	
evapor in soils, effect of temperature on	141-172	notch—continued.	
ement of.....	459-463	submerged, flow through.....	1101-1106
late and effect of arsenic on.....	i	triangular, flow through.....	1083-1088
city publication, announcement of.....	i	Wentworth, E. N., and Aubel, C. E. (paper),	
J. R., and Hubert, E. E. (paper), A		Inheritance of Fertility in Swine.....	1145-1160
nous Disease in Forest Nurseries Caused		Wheat. See <i>Triticum</i> .	
<i>Pendermium filamentosum</i>	781-785	Wilson, views on perpetuation of <i>Phytophthora</i>	
r—		<i>infestans</i>	97
α—		Wilt, bacterial, of cucurbits, dissemination	
effect of—		of.....	257-260
shape on discharge of.....	1129	Withers, W. A., and Carruth, F. E. (paper)	
size on discharge of.....	1129	Gossypol, the Toxic Substance in Cotton-	
discharge formula for.....	1135-1136	seed Meal.....	261-288
er irrigation.....	1127-1143	Wolf, F. A. (paper), Further Studies on Pea-	
advantages of.....	1143	nut Leafspot.....	891-902
construction of.....	1141-1143	Woodworth, C. W. (paper), A New Spray	
discharge from.....	1137-1139	Nozzle.....	1177-1182
experimental and computed discharges		Zea mays—	
from.....	1139-1141	absorption of boron by.....	584
use of.....	1141-1143	composition of seeds and seedlings of.....	456-457, 1162
αch—		effect of—	
Cipolletti, flow through.....	1073-1082	alkali salts on growth of.....	23ff
circular, flow through.....	1088	green manure on germination of.....	1162, 1165-1166
edges, requirements of, for free flow.....	1088-1090	translocation of mineral constituents of... 45	457
flow through.....	1051-1113	occurrence of manganese in.....	353
effect of contractions upon.....	1091-1098	resistance of, to alkali.....	22-24
measurement of head in.....	1090-1091	Zinnmeister, C. L., and Rosenbaum, J.	
relation of notch length to.....	1098-1101	(paper), <i>Alternaria panax</i> , the Cause of a	
rectangular, flow through.....	1063-1073	Rootrot of Ginseng.....	181-182

ADDITIONAL COPIES
OF THIS PUBLICATION MAY BE PROCURED FROM
THE SUPERINTENDENT OF DOCUMENTS
GOVERNMENT PRINTING OFFICE
WASHINGTON, D. C.
AT
10 CENTS PER COPY
SUBSCRIPTION PRICE, \$3.00 PER YEAR

▽

